

Predictive Regression and Robust Hypothesis Testing: Predictability Hidden by Anomalous Observations*

Lorenzo Camponovo

University of Lugano and University of St.Gallen

Olivier Scaillet

University of Geneva and Swiss Finance Institute

Fabio Trojani

University of Lugano and Swiss Finance Institute

First Version: November 2007; This Version: June 10, 2012

*We acknowledge the financial support of the Swiss National Science Foundation (NCCR FINRISK, grants 101312-103781/1, 100012-105745/1, and PDFM1-114533). We thank participants at the International Conference on Computational Management Science 2009 in Geneva, the International Conference on Robust Statistics 2009 in Parma, the International Conference on Computational and Financial Econometrics 2009 in Limassol, the Prospectus Workshop in Econometrics 2010 at Yale University, the seminar at the University of Ghent March 2010, the Conference on Resampling Methods and High Dimensional Data 2010 at Texas A&M University, ECARES seminar at ULB 2010 in Bruxelles, and seminars at the Universities of Orleans February 2012, the University of Frankfurt and University Bocconi May 2012, for helpful comments. Part of this research was performed when the first author was visiting Yale University as a post doctoral research fellow. Correspondence address: Fabio Trojani, Faculty of Economics, University of Lugano, Via Bu 13, CH-6900 Lugano, e-mail: Fabio.Trojani@usi.ch.

Abstract

Testing procedures for predictive regressions with lagged autoregressive variables imply a suboptimal inference in presence of small violations of ideal assumptions. We propose a novel testing framework resistant to such violations, which is consistent with nearly integrated regressors and applicable to multi-predictor settings, when the data may only approximately follow a predictive regression model. The Monte Carlo evidence demonstrates large improvements of our approach, while the empirical analysis produces a strong robust evidence of market return predictability, using predictive variables such as the dividend yield, the volatility risk premium or labor income.

Keywords: Predictive Regression, Stock Return Predictability, Bootstrap, Subsampling, Robustness.

JEL: C12, C13, G1.

1 Introduction

A large literature has investigated whether stock returns can be predicted by economic variables such as, e.g., the price-dividend ratio, proxies of labour income, or the interest rate.¹ The econometric approach to test for predictability is mostly based on a predictive regression of stock returns onto a set of lagged financial variables; see, e.g., Stambaugh (1999). Important differences between testing approaches in the literature arise because of the different test statistics, asymptotic theories or resampling approaches used to test the null hypothesis of no predictability. These differences lead in a number of cases to diverging results and conclusions.

Mankiw and Shapiro (1986) and Stambaugh (1986) note that in a setting with endogenous predictor and correlated innovations standard asymptotic theory causes small sample biases that may imply an overrejection of the hypothesis of no predictability. To mitigate the problem, recent studies propose tests based on bias-corrected estimators of predictive regressions. For instance, Stambaugh (1999), and Amihud, Hurvich and Wang (2008) introduce bias-corrected OLS estimators for the univariate and the multi-predictor setting, respectively.

Recent work has also considered the issue of endogenous integrated or nearly integrated predictors, following the evidence in Torous, Valkanov and Yan (2004) that various variables assumed to predict stock returns follow a local-to-unit root autoregressive process. Lewellen (2004), Torous, Valkanov and Yan (2004), and Campbell and Yogo (2006) introduce new testing procedures and more accurate unit-root and local-to-unit root asymptotics for predictive regression models with a single persistent predictor and correlated innovations.

A general approach to obtain tests that are less susceptible to finite sample biases or assumptions on the form of their asymptotic distribution can rely on nonparametric Monte Carlo simulation methods, such as the bootstrap or the subsampling. Ang and Bekaert (2007) use the bootstrap to quantify the bias of parameter estimation in a regression of stock returns on the lagged dividend yield and the interest rate. In a multi-predictor setting with nearly integrated

¹See Roze (1984), Fama and French (1988), Campbell and Shiller (1988), Nelson and Kim (1993), Goetzmann and Jorion (1995), Kothari and Shanken (1997), Campbell and Yogo (2006), Jansson and Moreira (2006), Polk, Thompson and Vuolteenaho (2006), Santos and Veronesi (2006), Bollerslev, Tauchen and Zhou (2009), among others.

regressors, Amihud, Hurvich and Wang (2008) compare the results of bootstrap tests to bias-corrected procedures and find the latter to have accurate size and good power properties. Wolf (2000) introduces subsampling tests of stock return predictability in single-predictor models.

As shown in Hall and Horowitz (1996) and Andrews (2002), among others, a desirable property of bootstrap tests is that they may provide asymptotic refinements of the sampling distribution of standard t -test statistics for testing the hypothesis of no predictability.² Moreover, as shown in Romano and Wolf (2001), Choi and Chue (2007) and Andrews and Guggenberger (2010), subsampling methods can produce reliable inference also in predictive regressions models with multiple nearly-integrated predictors.

A common feature of all above approaches to test predictability hypotheses is their reliance on procedures that can be heavily influenced by a small fraction of anomalous observations in the data. For standard OLS estimators and t -test statistics, this problem is well-known since a long time; see, e.g., Huber (1981) for a review. More recent research has also shown that inference provided by bootstrap and subsampling tests may be easily inflated by a small fraction of anomalous observations.³

Intuitively, this feature is due to the too high fraction of anomalous observations that is often simulated by standard bootstrap and subsampling procedures, when compared to the actual fraction of outliers in the original data. Unfortunately, this problem cannot be mitigated simply by applying standard bootstrap or subsampling methods to more robust statistics, so that robust bootstrap and subsampling schemes for time series are needed in order to develop more resistant tests for predictive regressions.

In this paper, we propose a new general method for hypothesis testing in predictive regressions with small samples and multiple nearly-integrated regressors, which is resistant to anomalous observations and other small violations of ideal assumptions in predictive regression

²In the sense that the errors made in approximating the true finite-sample distribution of the t -test statistic are of lower order with respect to the sample size than those implied by the conventional asymptotics.

³Following Huber seminal work, several authors have emphasized the potentially weak robustness features of many standard asymptotic testing procedures; see Heritier and Ronchetti (1994), Ronchetti and Trojani (2001), Mancini, Ronchetti and Trojani (2005), and Gagliardini, Trojani and Urga (2005), among others. Singh (1998), Salibian-Barrera and Zamar (2002), and Camponovo, Scaillet and Trojani (2012), among others, highlight the failing robustness of the inference implied by bootstrap and subsampling tests.

models. Our testing approach is based on a new class of robust resampling methodologies for time series, which ensure reliable inference properties in presence of anomalous observations. An additional desirable feature of our approach is that it simultaneously allows the identification of problematic observations, which may produce a weak performance of standard testing procedures. The more detailed contributions to the literature are as follows.

First, using Monte Carlo simulations we find that conventional hypothesis testing methods for predictive regressions, including bias-corrected tests, tests implied by near-to-unity asymptotics, and conventional bootstrap or subsampling tests, are dramatically non-resistant to even small fractions of anomalous observations in the data. Even though the test probability of rejecting a null by chance alone features some degree of resistance in our Monte Carlo experiments, the test ability to reject the null of no predictability when it is violated is in most cases drastically reduced.

Second, we quantify theoretically the robustness properties of bootstrap and subsampling tests of predictability in a time series context, borrowing from the concept of breakdown point, which is a measure of the degree of resistance of a testing procedure to outliers; see, e.g., Hampel (1971), Donoho and Huber (1983), and Hampel, Ronchetti, Rousseeuw and Stahel (1986). Our theoretical results confirm the dramatic non-robustness of conventional tests of predictability detected in our Monte Carlo findings.

Third, we develop a novel class of bootstrap and subsampling tests for times series, which are resistant to anomalous observations and are applicable in linear and nonlinear model settings at sustainable computational costs.⁴ We make use of our theory to propose a novel class of robust tests of predictability. We confirm by Monte Carlo simulations that these tests successfully limit the damaging effect of outliers, by preserving desirable finite sample properties in presence of anomalous observations.

Finally, we provide a robust analysis of the recent empirical evidence on stock return pre-

⁴Our robust resampling approach relies on the fast resampling idea put forward, among others, in Shao and Tu (1995), Davidson and McKinnon (1999), Hu and Kalbfleisch (2000), Andrews (2002), Salibián-Barrera and Zamar (2002), Gonçalves and White (2004), Hong and Scaillet (2006), Salibián-Barrera, Van Aelst and Willems (2006, 2007), and Camponovo, Scaillet and Trojani (2012). The methodology is applicable to a wide set of bootstrap and subsampling simulation schemes in the literature.

dictability for US stock market data. We study single-predictor and multi-predictor models, using several well-known predictive variables suggested in the literature, such as the lagged price-dividend ratio, the difference between option-implied volatility and realized volatility (Bollerslev, Tauchen and Zhou, 2009), and the share of labor income to consumption (Santos and Veronesi, 2006). Our robust tests of predictability produce the following novel empirical evidence.

First, we find that the dividend yield is a robust predictive variable of market returns, which is significant at the 5% significance in all our regressions, for each subperiod, sampling frequency and forecasting horizon considered. In univariate regressions with monthly data, the lagged dividend yield is significant at the 5% level according to the robust tests, in each window of 180 monthly observations from January 1980 to December 2010. In contrast, bias-corrected methods, near-to-unity asymptotics and standard resampling tests produce a weaker and more ambiguous evidence overall, e.g., by not rejecting the null of no predictability at the 10% significance level in the subperiod from January 1994 to December 2009 (bias-corrected methods and near-to-unity asymptotics) or in the subperiod from January 1995 to December 2010 (resampling tests). The significant predictive power of the dividend yield is confirmed by multi-predictor regressions including variance risk premium and labor income proxies. While the dividend yield is again significant at the 5% level in all cases using the robust tests, it is significant only at the 10% level using the conventional tests in the sample period 1994-2000, within monthly predictive regressions including the difference of implied and realized volatility as a predictive variable. It is not significant using conventional tests in the sample period 1955-2010 within quarterly predictive regression based on the share of labor income to consumption.

Second, we find that the difference between option-implied volatility and realized volatility is a robust predictive variable of future market returns at quarterly forecasting horizons. It is always significant at the 5% significance level in each window of 180 observations, using both robust and nonrobust testing approaches. This finding supports the remarkable market return forecasting ability of the variance risk premium, noted in Bollerslev, Tauchen and Zhou (2009) and confirmed in Bollerslev, Marrone and Zhou (2012) in an international context.

Third, using conventional testing approaches, we find that the evidence of return predictability associated with the ratio of labor income to consumption is either absent or weak in the sample periods 1955-2000 and 1965-2010, respectively. In contrast, the null of no predictability is always rejected at the 5% significance level by our robust testing method, indicating that the weak and ambiguous evidence produced by the conventional tests is likely a consequence of their low power in presence of anomalous observations.

Fourth, we exploit the properties of our robust testing method to identify potential anomalous observations that might excessively influence the diverging conclusions of conventional testing approaches. We find a fraction of less than about 5% of anomalous observations in the data, which tend to be more frequent during the NASDAQ bubble and the more recent financial crisis. Such influential data points, including the Lehman Brothers default on September 2008, the terrorist attack of September 2001, the Black Monday on October 1987 and the Dot-Com bubble collapse in August 2002, are largely responsible for the failure of conventional testing methods in uncovering the hidden predictability structures.

The rest of the paper is organized as follows. In Section 2, we introduce the usual predictive regression model, and we illustrate by simulation the robustness problem of some of the recent tests of predictability proposed in the literature. In Section 3, we study theoretically the robustness properties of tests based on resampling procedures in general time series settings. In Section 4, we introduce our robust approach and develop robust bootstrap and subsampling tests of predictability. In Section 5, we apply our robust testing procedure to US equity data and reconsider some of the recent empirical evidence on market return predictability. Section 6 concludes.

2 Predictability and Anomalous Observations

In this section, we introduce the benchmark predictive regression model and a number of recent testing methods proposed for testing the predictability of stock returns. Through Monte Carlo simulations, we study the finite-sample properties of these testing procedures both in presence and absence of anomalous observations. In Sections 2.1 and 2.2, we focus on bias-

corrected methods and testing procedures based on local-to-unity asymptotics. In Section 2.3, we consider more broadly testing approaches based on nonparametric Monte Carlo simulation methods, such as the bootstrap and the subsampling.

2.1 The Predictive Regression Model

We consider the predictive regression model:

$$y_t = \alpha + \beta x_{t-1} + u_t, \quad (1)$$

$$x_t = \gamma + \rho x_{t-1} + v_t, \quad (2)$$

where, for $t = 1, \dots, n$, y_t denotes the stock return at time t , and x_{t-1} is an economic variable observed at time $t - 1$, predicting y_t . The parameters $\alpha \in \mathbb{R}$ and $\beta \in \mathbb{R}$ are the unknown intercepts of the linear regression model and the autoregressive model, respectively, $\gamma \in \mathbb{R}$ is the unknown parameter of interest, $\rho \in \mathbb{R}$ is the unknown autoregressive coefficient, and $u_t \in \mathbb{R}$, $v_t \in \mathbb{R}$ are error terms with $u_t = v_t + e_t$, $e_t \in \mathbb{R}$, and e_t is a scalar random variable.

In this setting it is well-known that inference based on standard asymptotic theory suffers from small sample biases, which may imply an overrejection of the hypothesis of no predictability, $\mathcal{H}_0 : \gamma_0 = 0$, where γ_0 denotes the true value of the unknown parameter γ ; see Mankiw and Shapiro (1986), and Stambaugh (1986), among others. Moreover, as emphasized in Torous, Valkanov, and Yan (2004), various state variables considered as predictors in the model (1)-(2) are well approximated by a nearly integrated process. Consequently, this suggests a local-to-unity framework $\rho = 1 + c/n$, $c < 0$, for the autoregressive coefficient of model (2), which may imply a nonstandard asymptotic distribution for the OLS estimator $\hat{\gamma}_n$ of parameter γ .

Several recent testing procedures have been proposed in order to overcome these problems. Stambaugh (1999), Lewellen (2004), Amihud and Hurvich (2004), Polk, Thompson and Vuolteenaho (2006), and Amihud, Hurvich and Wang (2008, 2010), among others, propose bias-corrected procedures that correct the bias implied by the OLS estimator $\hat{\gamma}_n$ of parameter γ . Cavanagh, Elliott and Stock (1995), Torous, Valkanov and Yan (2004), and Campbell

and Yogo (2006), among others, introduce testing procedures based on local-to-unity asymptotics that provide more accurate approximations of the sampling distribution of the t -statistic $T_n = (\hat{\alpha}_n - \alpha_0) / \hat{\sigma}_n$ in nearly integrated settings, where $\hat{\sigma}_n$ is an estimate of the standard deviation of the OLS estimator $\hat{\alpha}_n$.

2.2 Bias Correction Methods and Near-to-Unity Asymptotic Tests

A common feature of bias-corrected methods and inference based on local-to-unity asymptotics is a non-resistance to anomalous observations, which may lead to conclusions determined by the particular features of a small subfraction of the data. Intuitively, this feature emerges because these approaches are based on statistical tools that can be sensitive to outliers or more general small deviations from the predictive regression model (1)-(2). Consequently, despite the good accuracy under the strict model assumptions, these testing procedures may become less efficient or biased even with a small fraction of anomalous observations in the data.

To illustrate the lack of robustness of this class of tests, we analyze through Monte Carlo simulation the bias-corrected method proposed in Amihud, Hurvich and Wang (2008) and the Bonferroni approach for the local-to-unity asymptotic theory introduced in Campbell and Yogo (2006). We first generate $N = 1,000$ samples $z_{(n)} = (z_1; \dots; z_n)$, where $z_t = (y_t; x_{t-1})'$, of size $n = 180$ according to model (1)-(2), with $v_t \sim N(0; 1)$, $e_t \sim N(0; 1)$, $\alpha = -1$, $\beta = 0$, $\rho = 0.9$, and $\alpha_0 \in [0; 0.15]$.⁵ In a second step, to study the robustness of the methods under investigation, we consider replacement outliers random samples $\tilde{z}_{(n)} = (\tilde{z}_1; \dots; \tilde{z}_n)$, where $\tilde{z}_t = (y_t; x_{t-1})'$ is generated according to,

$$y_t = (1 - p_t)y_t + p_t \cdot y_{3max}; \quad (3)$$

with $y_{3max} = 3 \cdot \max(y_1; \dots; y_n)$ and p_t is an iid $0 - 1$ random sequence, independent of process (1)-(2) such that $P[p_t = 1] = \alpha$. The probability of contamination by outliers is set to $\alpha = 4\%$, which is a small contamination of the original sample, compatible with the features of the real

⁵These parameter choices are in line with the Monte Carlo setting studied, e.g., in Choi and Chue (2007).

data set analyzed in the empirical study in Section 5.1.⁶

We study the finite sample properties of tests of the null hypothesis $\mathcal{H}_0 : \rho_0 = 0$ in the predictive regression model. For the different testing methods analyzed, Figure 1 plots the empirical frequency of rejection of null hypothesis \mathcal{H}_0 , with respect to different values of the alternative hypothesis $\rho_0 \in [0; 0.15]$. The nominal significance level of the test is 10%.

In the Monte Carlo simulation with non-contaminated samples (straight line), we find that the fraction of null hypothesis rejections of all procedures is quite close to the nominal level 10% when $\rho_0 = 0$. As expected, for increasing values of ρ_0 the power of the tests increases. For $\rho_0 = 0.1$, both methods have a frequency of rejection close to 70% and for $\rho_0 = 0.15$ a frequency larger than 95%. In the simulation with contaminated samples (dashed line), the size of all tests remains quite close to the nominal significance level. In contrast, the presence of anomalous observations dramatically deteriorates the power of both procedures. Indeed, for $\rho_0 > 0$, the frequency of rejection of the null hypothesis for both tests is much lower than in the non-contaminated case. The power of both tests is more flat and below 55% even for large values of ρ_0 . Unreported Monte Carlo results for different parameter choices and significance levels produce similar findings.

The results in Figure 1 highlight the lack of resistance to anomalous data of bias-corrected methods and inference based on local-to-unity asymptotics. Because of a small fraction of anomalous observations, the testing procedures become unreliable and are unable to reject the null hypothesis of no predictability, even for large values of ρ_0 . This is a relevant aspect for applications, in which typically the statistical evidence of predictability is weak.

To overcome this robustness problem, a natural approach is to develop more resistant versions of the nonrobust tests considered in our Monte Carlo exercise. However, this task may be hard to achieve in general.⁷ A more general approach to obtain tests that are less

⁶For the monthly data set in Section 5.1, the estimated fraction of anomalous observations in the sample period 1980-2010 is less than about 3.87%.

⁷To robustify the bias-corrected procedure in Amihud, Hurvich and Wang (2008), one would need to derive an expression for the bias of robust estimators of regressions, and then derive the asymptotic distribution of such bias-corrected robust estimators. For nearly-integrated settings, a robustification of the procedure proposed in Campbell and Yogo (2006) would require a not obvious extension of the robust local-to-unity asymptotics developed in Lucas (1995, 1997) for the predictive regression model.

susceptible to finite sample biases or assumptions on their asymptotic distribution can rely on nonparametric Monte Carlo simulation methods, such as the bootstrap or the subsampling. We address these methods in the sequel.

2.3 Bootstrap and Subsampling Tests

Nonparametric Monte Carlo simulation methods, such as the bootstrap and the subsampling, may provide improved inferences in predictive regression model (1)-(2) both in stationary or nearly integrated settings. As shown in Hall and Horowitz (1996) and Andrews (2002), for stationary data the block bootstrap may yield improved approximations to the sampling distribution of the standard t -statistics for testing predictability, having asymptotic errors of lower order in sample size. Moreover, as shown in Choi and Chue (2007) and Andrews and Guggenberger (2010), we can use the subsampling to produce correct inferences in nearly integrated settings. We first introduce in more detail block bootstrap and subsampling procedures. We then focus on predictive regression model (1)-(2) and study by Monte Carlo simulation the degree of resistance to anomalous observations of bootstrap and subsampling tests of predictability.

Consider a random sample $X_{(n)} = (X_1, \dots, X_n)$ from a time series of random vectors $X_i \in \mathbb{R}^{d_x}$, $d_x \geq 1$, and a general statistic $T_n := T(X_{(n)})$. Block bootstrap procedures split the original sample $X_{(n)}$ into overlapping blocks of size $m < n$. From these blocks, bootstrap samples $X_{(n)}^*$ of size n are randomly generated.⁸ Finally, the empirical distribution of statistic $T(X_{(n)}^*)$ is used to estimate the sampling distribution of $T(X_{(n)})$. Similarly, the more recent subsampling method applies statistic T directly to overlapping random blocks $X_{(m)}^*$ of size m strictly less than n .⁹ The empirical distribution of statistic $T(X_{(m)}^*)$ is then used to estimate the sampling distribution of $T(X_{(n)})$, under the assumption that the impact of the block size is asymptotically negligible ($m/n \rightarrow 0$).

In the predictive regression model (1)-(2), the usual t -test statistic for testing the null of no

⁸See, e.g., Hall (1985), Carlstein (1986), Künsch (1989) and Andrews (2004), among others. Alternatively, it is possible to construct the bootstrap samples using nonoverlapping blocks.

⁹See Politis, Romano and Wolf (1999), among others.

predictability is $T_n = (\hat{\alpha}_n - \alpha_0) = \hat{\alpha}_n$. Therefore, a block bootstrap test of the null hypothesis can be based on the block bootstrap statistic $T_{n,m}^{B*} = (\hat{\alpha}_{n,m}^{B*} - \hat{\alpha}_n) = \hat{\alpha}_{n,m}^{B*}$, where $\hat{\alpha}_{n,m}^{B*}$ is an estimate of the standard deviation of the OLS estimator $\hat{\alpha}_{n,m}^{B*}$ in a random bootstrap sample of size n , constructed using blocks of size m . Similarly, a subsampling test of the same null hypothesis can be based on the subsampling statistic $T_{n,m}^{S*} = (\hat{\alpha}_{n,m}^{S*} - \hat{\alpha}_n) = \hat{\alpha}_{n,m}^{S*}$, where $\hat{\alpha}_{n,m}^{S*}$ is now an estimator of the standard deviation of the OLS estimator $\hat{\alpha}_{n,m}^{S*}$ in a random overlapping block of size $m < n$.

It is well-known that OLS estimators and empirical averages are very sensitive to even small fractions of anomalous observations in the data; see, e.g., Huber (1981). Since block bootstrap and subsampling tests rely on such statistics, inference based on resampling methods may inherit the lack of robustness. To verify this intuition, we study through Monte Carlo simulation the finite-sample properties of block bootstrap and subsampling tests of predictability in presence of anomalous observations. For comparison, we consider the same simulation setting of the previous section and study tests of the null hypothesis $\mathcal{H}_0 : \alpha_0 = 0$, using symmetric (bootstrap and subsampling) confidence intervals for parameter α_0 .¹⁰

Figure 2 plots the empirical frequencies of rejection of null hypothesis \mathcal{H}_0 for different values of the alternative hypothesis $\alpha_0 \in [0; 0.15]$. The nominal significance level of the test is 10%, as before. With non-contaminated samples (straight line), we find that for all values of $\alpha_0 \in [0; 0.15]$ the frequency of rejection of block bootstrap and subsampling tests is close to the one of the bias-corrected method and the Bonferroni approach observed in the previous section. For $\alpha_0 = 0$, the size of the tests is close to the nominal level 10%, while for $\alpha_0 = 0.15$ the power is larger than 95%. Contaminations with anomalous observations strongly deteriorate the power of the tests. With contaminated samples (dashed line), the frequency of rejection of the null hypothesis is always less than 55%, even for large values of α_0 . In particular, when $\alpha_0 = 0.15$, the difference in power for the subsampling applied to non-contaminated and contaminated samples is larger than 50%.

¹⁰Appendix A shows how to construct symmetric confidence intervals for the parameter of interest, based on block bootstrap and subsampling distributions. For the selection of the block size m , we use the standard data-driven methods described in Appendix B.

The results in Figure 2 show that bootstrap and subsampling tests inherit, and to some extent exacerbate, the lack of robustness of OLS estimators for predictive regressions. To robustify the inference produced by resampling methods, a natural idea is to apply standard bootstrap and subsampling simulation schemes to a more robust statistic, such as, e.g., a robust estimator of linear regression. Unfortunately, as shown in Singh (1998), Salibián-Barrera and Zamar (2002), and Camponovo, Scaillet and Trojani (2012) for iid settings, resampling a robust statistic does not yield a robust inference, because standard block bootstrap and subsampling procedures have an intrinsic non-resistance to outliers. Intuitively, this problem arises because the fraction of anomalous observations generated in bootstrap and subsampling blocks is often much higher than the fraction of outliers in the data. To solve this problem, it is necessary to analyze more systematically the robustness of bootstrap and subsampling methods for time series.

3 Robust Resampling Inference and Quantile Breakdown Point

We characterize theoretically the robustness of bootstrap and subsampling tests in predictive regression settings. Section 3.1 introduces the notion of a quantile breakdown point, which is a measure of the global resistance of a resampling method to anomalous observations. Section 3.2 quantifies and illustrates the quantile breakdown point of standard bootstrap and subsampling tests in predictive regression models. Finally, Section 3.3 derives explicit bounds for quantile breakdown points, which quantify the degree of resistance to outliers of bootstrap and subsampling tests for predictability, before applying them to the data.

3.1 Quantile Breakdown Point

Given a random sample $X_{(n)}$ from a sequence of random vectors $X_i \in \mathbb{R}^{d_x}$, $d_x \geq 1$, let $X_{(n,m)}^{B*} = (X_1^* \dots X_n^*)$ denote a block bootstrap sample, constructed using overlapping blocks of size m . Similarly, let $X_{(n,m)}^{S*} = (X_1^* \dots X_m^*)$ denote an overlapping subsampling block. We denote by

$T_{n,m}^{K*} := T(X_{(n,m)}^{K*})$, $K = S; B$, the corresponding block bootstrap and subsampling statistics, respectively.¹¹ For $t \in (0; 1)$, the quantile $Q_{t,n,m}^{K*}$ of $T_{n,m}^{K*}$ is defined by

$$Q_{t,n,m}^{K*} = \inf\{x | P^*(T_{n,m}^{K*} \leq x) \geq t\}; \quad (4)$$

where P^* is the probability measure induced by the block bootstrap or the subsampling method and, by definition, $\inf(\emptyset) = \infty$.

Quantile $Q_{t,n,m}^{K*}$ is effectively a useful nonparametric estimator of the corresponding finite-sample quantile of statistic $T(X_1; \dots; X_n)$. We study the breakdown point of quantile (4), i.e., the smallest fraction of outliers in the original sample such that $Q_{t,n,m}^{K*}$ diverges to infinity. Intuitively, when a breakdown occurs, inference about the distribution of $T(X_1; \dots; X_n)$ based on bootstrap or subsampling tests becomes meaningless. Estimated test critical values may be arbitrarily large and confidence intervals be arbitrarily wide. In these cases, the size and power of bootstrap and subsampling tests can collapse to zero or one in presence of anomalous observations, making these inference procedures useless.

We make use of the concept of quantile breakdown point, in order to characterize the largest fraction of anomalous observations in the data, such that bootstrap and subsampling inferences can still provide some reliable information in applications. Borrowing the notation in Genton and Lucas (2003), we formally define the breakdown point of the t -quantile $Q_{t,n,m}^{K*} := Q_{t,n,m}^{K*}(X_{(n)})$ as,

$$b_{t,n,m}^K := \frac{1}{n} \cdot \left[\inf_{\{1 \leq p \leq \lceil n/2 \rceil\}} \{p | \text{there exists } Z_{n,p}^\zeta \in \mathcal{Z}_{n,p}^\zeta \text{ such that } Q_{t,n,m}^{K*}(X_{(n)} + Z_{n,p}^\zeta) = +\infty\} \right]; \quad (5)$$

where $\lceil x \rceil = \inf\{n \in \mathbb{N} | x \leq n\}$, and $\mathcal{Z}_{n,p}^\zeta$ denotes the set of all n -samples $Z_{n,p}^\zeta$ with exactly p non-zero components that are d_x -dimensional outliers of size $\zeta \in \mathbb{R}^{d_x}$.¹²

¹¹We focus for brevity on one-dimensional real-valued statistics. However, as discussed for instance in Singh (1998) in the iid context, our results for time series can be naturally extended to multivariate and scale statistics.

¹²When $p > 1$, we do not necessarily assume outliers ζ_1, \dots, ζ_p to be all equal to ζ , but we rather assume existence of constants c_1, \dots, c_p , such that $\zeta_i = c_i \zeta$. To better capture the presence of outliers in predictive regression models, our definitions for the breakdown point and the set $\mathcal{Z}_{n,p}^\zeta$ of all n -components outlier samples are slightly different from those proposed in Genton and Lucas (2003) for general settings. However, the results

Literally, $b_{t,n,m}^K$ is the smallest fraction of anomalous observations of arbitrary size, in a generic outlier-contaminated sample $X_{(n)} + Z_{n,p}^\zeta$, such a quantile $Q_{t,n,m}^{K*}$, estimated by a bootstrap or a subsampling Monte Carlo simulation scheme, can become meaningless. Therefore, quantifying $b_{t,n,m}^K$ in general for bootstrap and subsampling tests of predictability, in dependence of the statistics and testing approaches used, is key in order to understand which approaches ensure some resistance to anomalous observations and which do not, even before looking at the data.

3.2 Quantile Breakdown Point and Predictive Regression

The quantile breakdown point of conventional block bootstrap and subsampling tests for predictability in Section 2.3 depends directly on the breakdown properties of OLS estimator $\hat{\alpha}_n$. The breakdown point b of a statistics $T_n = T(X_{(n)})$ is simply the smallest fraction of outliers in a contaminated sample $X_{(n)} + Z_{n,nb}^\zeta$ such that $T(X_{(n)} + Z_{n,nb}^\zeta) = \pm\infty$ for some outlier-contaminated sample $Z_{n,nb}^\zeta \in \mathcal{Z}_{n,nb}^\zeta$. b is explicitly known in some cases and it can be gauged most of the time, for instance by means of simulations and sensitivity analysis. Most non-robust statistics, like OLS estimators for linear regression, have a breakdown point $b = 1/n$. Therefore, the breakdown point of conventional block bootstrap and subsampling quantiles in predictive regression settings also equals $1/n$. In other words, a single anomalous observation in the original data is sufficient to produce a meaningless inference implied by bootstrap or subsampling quantiles in standard tests of predictability.

It is straightforward to illustrate these features in a Monte Carlo simulation that quantifies the sensitivity of block bootstrap and subsampling quantiles to data contaminations by a single outlier, where the size of the outlier is increasing. We first simulate $N = 1,000$ random samples $Z_{(n)} = (z_1, \dots, z_n)$ of size $n = 120$, where $z_t = (y_t, x_{t-1})'$ follows model (1)-(2), $v_t \sim N(0;1)$, $e_t \sim N(0;1)$, $\alpha = -1$, $\beta = 0$, $\gamma = 0.9$, and $\alpha_0 = 0$. For each Monte Carlo sample, we define in our paper can be easily modified to cover alternative definitions of breakdown point and outlier sets $\mathcal{Z}_{n,p}^\zeta$.

in a second step

$$y_{\max} = \arg \max_{y_1, \dots, y_n} \{w(y_i) | w(y_i) = y_i - \beta_0 x_{i-1}, \text{ under } \mathcal{H}_0 : \beta_0 = 0\} ; \quad (6)$$

and we modify y_{\max} over the interval $[y_{\max}; y_{\max} + 5]$. This means that we contaminate the predictability relationship by an anomalous observation for only one single data point in the full sample. We study the sensitivity of the Monte Carlo average length of confidence intervals for parameter β_0 , estimated by the standard block bootstrap and the subsampling. This is a natural exercise, as the length of the confidence interval for parameter β_0 is in a one-to-one relation with the critical value of the test of the null of no predictability ($\mathcal{H}_0: \beta_0 = 0$). For the sake of comparison, we also consider confidence intervals implied by the bias-corrected testing method in Amihud, Hurvich and Wang (2008) and the Bonferroni approach proposed in Campbell and Yogo (2006).

For all tests under investigation, Figure 4 and 5 plot the relative increase of the average confidence interval length in our Monte Carlo simulations, under contamination by a single outlier of increasing size. We find that all sensitivities are basically linear in the size of the outlier, confirming that a single anomalous observation can have an arbitrarily large impact on the critical values of those tests and make the test results potentially useless, as implied by their quantile breakdown point of $1/n$.

3.3 Quantile Breakdown Point Bounds

To obtain bootstrap and subsampling tests with more favorable breakdown properties, it is necessary to apply resampling procedures to a robust statistic with nontrivial breakdown point ($b > 1/n$), such as, e.g., a robust estimator of linear regression. Without loss of generality, let $T_n = T(X_{(n)})$ be a statistic with breakdown point $1/n < b \leq 0.5$.

We compute explicit quantile breakdown point bounds, which characterize the resistance of bootstrap and subsampling tests to anomalous observations, in dependence of relevant parameters, such as n , m , t , and b .¹³ These results are the starting point to develop more resistant

¹³Similar results can be obtained for the subsampling and the block bootstrap based on nonoverlapping

subsampling and bootstrap tests with a large of resistance to anomalous observations.

Theorem 1 *Let b be the breakdown point of T_n and $t \in (0;1)$. The quantile breakdown point $b_{t,n,m}^S$ and $b_{t,n,m}^B$ of subsampling and block bootstrap procedures, respectively, satisfy following properties,*

$$\frac{\lceil mb \rceil}{n} \leq b_{t,n,m}^S \leq \frac{1}{n} \cdot \left[\inf_{\{p \in \mathbb{N}, p \leq r-1\}} \left\{ p \cdot \lceil mb \rceil \mid p > \frac{(1-t)(n-m+1) + \lceil mb \rceil - 1}{m} \right\} \right]; \quad (7)$$

$$\frac{\lceil mb \rceil}{n} \leq b_{t,n,m}^B \leq \frac{1}{n} \cdot \left[\inf_{\{p_1, p_2\}} \left\{ p = p_1 \cdot p_2 \mid P \left(\text{BIN} \left(r; \frac{mp_2 - p_1 + 1}{n - m + 1} \right) \geq \left\lceil \frac{nb}{p_1} \right\rceil \right) > 1 - t \right\} \right] \quad (8)$$

where $p_1, p_2 \in \mathbb{N}$, with $p_1 \leq m; p_2 \leq r - 1$.

We illustrate the implications of Theorem 1 by computing in Table 1 lower and upper bounds for the breakdown point of subsampling and bootstrap quantiles, using a sample size $n = 120$ and a maximal statistic breakdown point ($b = 0.5$).¹⁴ We find that even for a highly robust statistic with maximal breakdown point ($b = 0.5$), the subsampling implies a very low quantile breakdown point, which increases with the block size but is also very far from the maximal value $b = 0.5$. For instance, for a block size $m = 10$, the 0.95-quantile breakdown point is between 0.0417 and 0.0833. In other words, even though a statistic is resistant to large fractions of anomalous observations, the implied subsampling quantile can collapse with just 5 outliers out of 100 observations.¹⁵ Similar results arise for the bootstrap quantiles. Even though the bounds are less sharp than for the subsampling, quantile breakdown points are again clearly smaller than the breakdown point of the statistic used.¹⁶

blocks. The results for the block bootstrap can also be modified to cover asymptotically equivalent variations, such as the stationary bootstrap of Politis and Romano (1994).

¹⁴The term $\frac{(1-t)(n-m+1)}{m}$ represents the number of degenerated subsampling statistics necessary in order to cause the breakdown of $Q_{t,n,m}^{S*}$, while $\frac{\lceil mb \rceil}{n}$ is the fraction of outliers which is sufficient to cause the breakdown of statistic T in a block of size m . Note that the breakdown point formula for the iid bootstrap in Singh (1998) emerges as a special case of the formula (8), for $m = 1$.

¹⁵This breakdown point is also clearly lower than in the iid case; see Camponovo, Scaillet and Trojani (2012). For instance, for $m = 10$, the 0.95-quantile breakdown point of the overlapping subsampling is 0.23 in iid settings. Since in a time series setting the number of possible subsampling blocks of size m is typically lower than the number of iid subsamples of size m , the breakdown of a statistic in one random block tends to have a larger impact on the subsampling quantile than in the iid case.

¹⁶These quantile breakdown point bounds are again clearly lower than in the iid setting. For instance, for

Overall, the results in Theorem 1 imply that subsampling and bootstrap tests for time series feature an intrinsic non-resistance to anomalous observations, which cannot be avoided, simply by applying conventional resampling approaches to more robust statistics.

4 Robust Bootstrap and Subsampling

When using a highly robust statistic with large breakdown point, the bootstrap and the subsampling still imply an important non-resistance to anomalous observations, which is consistent with our Monte Carlo results in the predictive regression model. To overcome the problem, it is necessary to introduce a novel class of more robust bootstrap and subsampling tests in the time series context.¹⁷ Section 4.1 introduces our robust bootstrap and subsampling approach and Section 4.2 demonstrates its favorable breakdown properties. Section 4.3 applies the theory to introduce new robust bootstrap and subsampling tests for predictability in predictive regression models.

4.1 Definition

Given the original sample $X_{(n)} = (X_1, \dots, X_n)$, we consider the class of robust M-estimators $\hat{\theta}_n$ for parameter $\theta \in \mathbb{R}^d$, defined as the solution of the estimating equations

$$\psi_n(X_{(n)}; \hat{\theta}_n) := \frac{1}{n} \sum_{i=1}^n g(X_i; \hat{\theta}_n) = 0; \quad (9)$$

where $\psi_n(X_{(n)}; \cdot) : \mathbb{R}^d \rightarrow \mathbb{R}^d$ depends on parameter θ and a bounded estimating function g . Boundedness of function g is a characterizing feature of robust M-estimators, such as, e.g., robust estimators of linear regression; see Hampel, Ronchetti, Rousseeuw and Stahel (1986) for a review.

$m = 30$, the 0.95-quantile breakdown point for time series is less than 0.25, but it is 0.425 for iid settings, from the results in Camponovo, Scaillet and Trojani (2012).

¹⁷We develop such robust methods for the broader class of M-estimators, borrowing from the fast resampling approach considered, among others, in Shao and Tu (1995), Davidson and McKinnon (1999), Hu and Kalbfleisch (2000), Andrews (2002), Salibián-Barrera and Zamar (2002), Gonçalves and White (2004), Hong and Scaillet (2006), Salibián-Barrera, Van Aelst and Willems (2006, 2007), and Camponovo, Scaillet and Trojani (2012).

Conventional bootstrap (subsampling) methods solve equation $k(X_{(n,m)}^{K*}; \hat{\theta}_{n,m}^{K*}) = 0$, for each bootstrap (subsampling) random sample $X_{(n,m)}^{K*}$, which can be a computationally demanding task. Instead, we consider a standard Taylor expansion of (9) around the true parameter θ_0 ,

$$\hat{\theta}_n - \theta_0 = -[\nabla_{\theta} \ell_n(X_{(n)}; \theta_0)]^{-1} \ell_n(X_{(n)}; \theta_0) + o_p(1); \quad (10)$$

where $\nabla_{\theta} \ell_n(X_{(n)}; \theta_0)$ is the derivative of function ℓ_n with respect to parameter θ . Based on this expansion, we can use $-[\nabla_{\theta} \ell_n(X_{(n)}; \hat{\theta}_n)]^{-1} k(X_{(n,m)}^{K*}; \hat{\theta}_n)$ as an approximation of $\hat{\theta}_{n,m}^{K*} - \hat{\theta}_n$ in the definition of the resampling scheme estimating the sampling distribution of $\hat{\theta}_n - \theta_0$. This approach avoids computing $\hat{\theta}_{n,m}^{K*}$ and $[\nabla_{\theta} k(X_{(n,m)}^{K*}; \hat{\theta}_n)]^{-1}$ in each bootstrap or subsampling sample, which is a markable computational advantage that produces a fast numerical procedure. This is an important improvement over conventional resampling schemes, which can easily become unfeasible when applied to robust statistics.

Definition 2 Given a normalization constant τ_n such that $\tau_n \rightarrow \infty$ as $n \rightarrow \infty$, a robust fast resampling distribution for $\tau_n(\hat{\theta}_n - \theta_0)$ is defined by

$$L_{n,m}^{K*}(x) = \frac{1}{N} \sum_{s=1}^N \mathbb{I} \left(\tau_n \left(-[\nabla_{\theta} \ell_n(X_{(n)}; \hat{\theta}_n)]^{-1} k(X_{(n,m),s}^{K*}; \hat{\theta}_n) \right) \leq x \right); \quad (11)$$

where $\mathbb{I}(\cdot)$ is the indicator function and s indexes the N possible random samples generated by the bootstrap and subsampling procedures, respectively.

General assumptions under which (11) consistently estimates the unknown asymptotic distribution of $\tau_n(\hat{\theta}_n - \theta_0)$ in a time series context are given, e.g., in Hong and Scaillet (2006) for the subsampling (Assumption 1) and in Goncalves and White (2004) for the bootstrap (Assumption A and Assumptions 2.1 and 2.2).¹⁸

¹⁸Let $L(\cdot)$ be the limit distribution function of $\tau_n(\hat{\theta}_n - \theta_0)$. Under regularity conditions, it follows, for any continuity point of $L(\cdot)$: $L_{n,m}^{K*}(x) \rightarrow L(x)$ in probability, as $N, n \rightarrow \infty$, and additionally $m \rightarrow \infty$, $m/n \rightarrow 0$ in the subsampling case. If $L(\cdot)$ is continuous, then: $\sup_x |L_{n,m}^{K*}(x) - L(x)| \rightarrow 0$ in probability. Moreover, as noted, e.g., in Hall, Horowitz and Jing (1995), several standard resampling methods imply the distortion $E^*[\hat{\theta}_{n,m}^{K*}] \neq E^*[\hat{\theta}_n]$, where $E^*[\cdot]$ denotes the expectation under the bootstrap or the subsampling distribution P^* , which decreases the rate of convergence. A solution to this problem can be based on the recentered statistic

4.2 Robust Resampling Methods and Quantile Breakdown Point

In the computation of the resampling distribution (11), we only need consistent point estimates for parameter vector θ_0 and matrix $-\nabla_{\theta} \psi_n(X_{(n)}; \theta_0)^{-1}$, based on the original sample $X_{(n)}$. These estimates are simply given by $\hat{\theta}_n$ and $-\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)^{-1}$, respectively.

A closer inspection of quantity $-\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)^{-1} k(X_{(n,m),s}^{K*}; \hat{\theta}_n)$ in Definition 2 reveals important implications for the breakdown properties of the robust fast resampling distribution (11). Indeed, this quantity can degenerate only when either (i) matrix $\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)$ is singular or (ii) estimating function g is not bounded. However, since we are making use of a robust (bounded) estimating function g , situation (ii) cannot arise. Therefore, one intuitively expects the breakdown of the quantiles of robust fast resampling distribution (11) to arise only when condition (i) is realized. These arguments imply the following immediate corollary.

Corollary 3 *Let b be the breakdown point of the robust M-estimator $\hat{\theta}_n$ defined in (9). The t -quantile breakdown point of resampling distribution (11) equals $b_{t,n,m}^K = \min(b; b_{\nabla\psi})$, where $b_{\nabla\psi}$ is the breakdown point of matrix $\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)$, defined by:*

$$b_{\nabla\psi} = \frac{1}{n} \cdot \inf_{1 \leq p \leq \lfloor n/2 \rfloor} \{p \mid \text{there exists } Z_{n,p}^\zeta \in \mathcal{Z}_{n,p}^\zeta \text{ such that } \det(\nabla_{\theta} \psi_n(X_{(n)} + Z_{n,p}^\zeta; \hat{\theta}_n)) = 0\}: \quad (12)$$

Since $b_{t,n,m}^K$ is the minimum of b and the breakdown point of matrix $\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)$, the quantile breakdown point of the robust fast resampling distribution in Definition 2 is maximal whenever

$$b_{\nabla\psi} \geq b$$

4.3 Robust Predictive Regression and Hypothesis Testing

We develop a new class of easily applicable robust bootstrap and subsampling tests for the null hypothesis of no predictability in predictive regression models. To this end, consider the predictive regression model

$$y_t = \beta' w_{t-1} + u_t ; t = 1, \dots, n ; \quad (13)$$

with $\beta = (\beta_0, \beta_1)'$ and $w_{t-1} = (1, x_{t-1})'$, and denote by $Z_{(n)} = (Z_1, \dots, Z_n)$ an observation sample generated according to (13), where $Z_t = (y_t, w_{t-1})'$.

According to Definition 2, a robust estimator of predictive regression is needed, featuring a nontrivial breakdown point $b > 1/n$ and a bounded estimating function g , in order to obtain robust bootstrap and subsampling tests with our approach. Several such estimators are available in the literature, which imply corresponding robust bootstrap and subsampling procedures. Among those estimators, a convenient choice is the standard Huber estimator $\hat{\beta}_n^R$ of regression, which ensures together good robustness properties and moderate computational costs.

Given a positive constant c , $\hat{\beta}_n^R$ is the M -estimator that solves the equation

$$n, c(Z_{(n)}; \hat{\beta}_n^R) := \frac{1}{n} \sum_{t=1}^n g_c(y_t; w_{t-1}; \hat{\beta}_n^R) = 0 ; \quad (14)$$

with estimating function g_c defined by

$$g_c(y_t; w_{t-1}; \beta) = (y_t - \beta' w_{t-1}) w_{t-1} \cdot \min \left(1, \frac{c}{\| (y_t - \beta' w_{t-1}) w_{t-1} \|} \right) ; \quad (15)$$

Constant $c > 0$ is useful in order to tune the degree of robustness of estimator $\hat{\beta}_n^R$ in relevant applications and can be determined in a fully data-driven way.²⁰ Note that, as required by our robust resampling approach, the norm of estimating function g_c in Equation (14) is bounded (by constant c), and the breakdown point of estimator $\hat{\beta}_n^R$ is maximal ($b = 0.5$).

²⁰Appendix B presents in detail the data-driven method that can be used for the selection of tuning constant c .

4.3.1 Nonstudentized Tests

By applying the robust fast approach in Definition 2 to the estimating function (14), we can estimate the sampling distribution of the nonstudentized statistic $T_n^{NS} = \sqrt{n}(\hat{\alpha}_n^R - \alpha_0)$, using the following robust fast resampling distribution:

$$L_{n,m}^{NS,K*}(x) = \frac{1}{N} \sum_{s=1}^N \mathbb{I} \left(\sqrt{k} \left(-[\nabla_{\theta_{n,c}}(Z_{(n)}; \hat{\alpha}_n)]^{-1} \nabla_{k,c}(Z_{(n,m),s}^{K*}; \hat{\alpha}_n) \right) \leq x \right) ; K = B; S; \quad (16)$$

where $\alpha_0 = (\alpha_0; \beta_0)'$ and $k = n$ for the block bootstrap ($k = m$ for the subsampling).

A key property of resampling distribution (16) is that it implies a maximal quantile breakdown point, i.e., the largest achievable degree of resistance to anomalous observations, independent of the probability level t and the selected block size m , both for the bootstrap and the subsampling. This feature follows directly from the fact that the breakdown point of the robust Huber estimator (14) is $b = 0.5$, while because $b_{\nabla\psi_c} \geq 0.5$ (see the proof in Appendix C).

Corollary 4 *Let $t \in (0;1)$. The t -quantile breakdown point of the resampling distribution (16) is $b_{t,n,m}^K = 0.5$, $K = B; S$.*

Using nonstudentized statistics, robust resampling distribution (16) provides consistent estimators of the sampling distribution of $T_n^{NS} = \sqrt{n}(\hat{\alpha}_n^R - \alpha_0)$ in stationary time series settings. Using studentized statistics, we can additionally obtain convergence in nonstationary settings and potentially also produce a faster convergence to the asymptotic distribution.

4.3.2 Studentized Tests

Consider the studentized statistic $T_n = \sqrt{n}[\hat{\alpha}_n^R]^{-1/2}(\hat{\alpha}_n^R - \alpha_0)$, where $\hat{\alpha}_n^R$ is an estimator of the asymptotic variance of $\hat{\alpha}_n^R$. Also in this case, we can use the robust fast approach in Definition 2 with minor modifications. More precisely, we propose to estimate the sampling distribution

of statistic T_n by the following robust fast resampling distribution:

$$L_{n,m}^{K*}(x) = \frac{1}{N} \sum_{s=1}^N \mathbb{I} \left(\sqrt{k} \left(- [\hat{\Lambda}_{k,s}^{R*}]^{-1/2} [\nabla_{\theta_{k,c}}(Z_{(n,m),s}^{K*})]^{-1} \right) \leq x \right); \quad (17)$$

where $k = n$ for the block bootstrap ($k = m$ for the subsampling) and $\hat{\Lambda}_{k,s}^{R*} = \hat{\Lambda}_n^R(Z_{(n,m),s}^{K*})$ is an estimate of the asymptotic variance of the robust M-estimator (14) based on the s -th block bootstrap (subsampling) random block.

The quantile breakdown point properties of resampling distribution (17) are more complex than those obtained in the unstudentized case. Unlike the exact results in Theorem 1, we provide the following bounds for the resulting breakdown point quantiles.

Theorem 5 *For simplicity, let $r = n=m \in \mathbb{N}$. The t -quantile breakdown points $b_{t,n,m}^B$ and $b_{t,n,m}^S$ of the robust block bootstrap and robust subsampling distributions (17), respectively, are given by*

$$b_{t,n,m}^S = \frac{1}{n} \left[\inf_{\{p \in \mathbb{N}, p \leq n-m+1\}} \left\{ m + p \mid p > (1-t)(n-m+1) - 1 \right\} \right]; \quad (18)$$

$$b_{t,n,m}^B = \frac{1}{n} \left[\inf_{\{p \in \mathbb{N}, p \leq n-m+1\}} \left\{ m + p \mid P \left(\text{BIN} \left(r; \frac{p+1}{n-m+1} \right) = r \right) > 1-t \right\} \right]; \quad (19)$$

where $\text{BIN}(N; q)$ is a Binomial random variable with parameters N and $q \in (0, 1)$.

Formulas (18) and (19) improve on the results in Equations (7) and (8) for the conventional subsampling and bootstrap, respectively. The quantile breakdown point of the robust block bootstrap and subsampling approach is often much higher than the one of conventional resampling methods. Table 2 quantifies these differences. For instance, for $m = 10; 20$, the 0.95-quantile of the robust block bootstrap is maximal. Similarly, the robust subsampling quantile breakdown points in Table 2 are considerably larger than those in Table 1 for conventional subsampling methods, even if they do not always attain the upper bound of 0.5.

In contrast to the unstudentized case, the quantile breakdown point of robust resampling distribution (17) is not always maximal, because we need to compute matrices $\hat{\Lambda}_k^{R*}$ and $\nabla_{\theta_{k,c}}(Z_{(n,m)}^{K*})$ in each bootstrap or subsampling block. While this approach can yield con-

sistency also in nonstationary settings and a potentially improved convergence, the additional estimation step can imply a loss in the resistance of the whole procedure to anomalous observations. Thus, a tradeoff arises between resistance to anomalous observations and improved finite-sample inference, which has to be considered and evaluated case-by-case in applications.

4.3.3 Monte Carlo Evidence

To quantify the implications of Corollary 4, we can study the sensitivity of confidence intervals estimated by the robust block bootstrap and the robust subsampling distributions (16), with respect to contaminations by anomalous observations of increasing size. To this end, we consider the same Monte Carlo setting of Section 3.2. We plot in Figure 6 the percentage increase of the length in the average estimated confidence interval, with respect to contaminations of the available data by a single anomalous observation of increasing size. In evident contrast to the findings for conventional bootstrap and subsampling tests, Figure 6 shows that the inference implied by our robust approach is largely insensitive to outliers, with a percentage increase in the average confidence interval length that is less than 1%, even for an outliers of size $y_{max} + 5$.

The robustness of resampling distribution (16) has favorable implications for the power of bootstrap and subsampling tests in presence of anomalous observations. For the same Monte Carlo setting of Sections 2.2 and 2.3, Figure 3 shows that in presence of non-contaminated samples (straight line) the frequencies of null hypothesis rejections of robust resampling approaches are again very close to those observed for nonrobust methods. For instance, the frequency of rejections when $\rho_0 = 0$ is close to the nominal level 10% of the test, while the power is larger than 95% for $\rho_0 = 0.15$. In presence of anomalous observations (dashed line), robust approaches still provide an accurate empirical size close to the actual nominal level, as well as a power curve that is close to the one obtained in the non-contaminated Monte Carlo simulation. In particular, in contrast to the standard tests, the power of both robust tests is above 90% for $\rho_0 = 0.15$.

Unreported Monte Carlo results for the robust block bootstrap and subsampling distributions of studentized tests (17) confirm the improved resistance properties of our approach.

5 Empirical Evidence of Return Predictability

Using our robust bootstrap and subsampling tests, we revisit from a robustness perspective the recent empirical evidence on return predictability for US stock market data. We study single-predictor and multi-predictor settings, using several well-known predictive variables suggested in the literature, such as the lagged dividend yield, the difference between option-implied volatility and realized volatility (Bollerslev, Tauchen and Zhou, 2009) and the share of labor income to consumption (Santos and Veronesi, 2006). We compare the evidence produced by our robust bootstrap and subsampling tests of predictability with the results of recent testing methods proposed in the literature, including the bias-corrected method in Amihud, Hurvich and Wang (2008), the Bonferroni approach for local-to-unity asymptotics in Campbell and Yogo (2006), and conventional bootstrap and subsampling tests.

The dividend yield is the most common predictor of future stock returns, as suggested by a simple present-value logic.²¹ However, its forecasting ability has been called into question, e.g., by the ambiguous empirical evidence of studies not rejecting the null of no predictability for a number of forecasting horizons and sample periods; see, e.g., Goyal and Welch (2003), and Ang and Bekaert (2007), among others. Whether these ambiguous results are related to the weakness of conventional tests in detecting predictability structures masked by anomalous observations, is an empirical question that we can analyze using our robust testing method.

The empirical study is articulated in three parts. Section 5.1 studies the forecast ability of the lagged dividend yield for explaining monthly S&P 500 index returns, in a predictive regression model with a single predictor. This study allows us to compare the results of our methodology with those of the Bonferroni approach for local-to-unity asymptotics, which is applicable to univariate regression settings. Section 5.2 considers instead models with two predictive variables. In Section 5.2.1, we test the predictive power of the dividend yield and the variance risk premium, for quarterly S&P 500 index returns sampled at a monthly frequency in periods marked by a financial bubble and a financial crisis. Finally, Section 5.2.2 tests

²¹See, e.g., Roze (1984), Campbell and Shiller (1988), Fama and French (1988), Stambaugh (1999), Lewellen (2004), Torous, Valkanov and Yan (2004), Lettau and Ludvigson (2005), and Campbell and Yogo (2006).

the predictive power of the dividend yield and the ratio of labor income to consumption for predicting quarterly value-weighted CRSP index returns.²²

5.1 Single-Predictor Model

We consider monthly S&P 500 index returns from Shiller (2000), $R_t = (P_t + d_t)/P_{t-1}$, where P_t is the end of month real stock price and d_t the real dividend paid during month t . Consistent with the literature, the annualized dividend series D_t is defined as

$$D_t = d_t + (1 + r_t)d_{t-1} + (1 + r_t)(1 + r_{t-1})d_{t-2} + \cdots + (1 + r_t) \cdots (1 + r_{t-10})d_{t-11}; \quad (20)$$

where r_t is the one-month maturity Treasury-bill rate. We estimate the predictive regression model

$$\ln(R_t) = \alpha + \beta \ln\left(\frac{D_{t-1}}{P_{t-1}}\right) + \varepsilon_t; \quad t = 1, \dots, n; \quad (21)$$

and test the null of no predictability, $\mathcal{H}_0 : \beta = 0$.

We collect monthly observations in the sample period 1980-2010 and estimate the predictive regression model using rolling windows of 180 observations. Figure 7 plots the 90%-confidence intervals for parameter β in the sample period 1980-2010, while Table 3 reports the detailed point estimates and test results for the different testing procedures in the four subperiods 1980-1995, 1985-2000, 1990-2005, 1995-2010.

We find that while the robust bootstrap and subsampling tests always clearly reject the hypothesis of no predictability at the 5%-significance level, the conventional testing approaches produce a weaker and more ambiguous predictability evidence. For instance, the bootstrap and subsampling tests cannot reject \mathcal{H}_0 at the 10% significance level in subperiod 1984-1999, while the bias-corrected method and the Bonferroni approach fail to reject \mathcal{H}_0 at the 10% significance level in the subperiod 1995-2010.

It is interesting to study to which extent anomalous observations in sample periods 1984-1999 and 1995-2010 might have caused the diverging conclusions of robust and nonrobust

²²We also consider regressions with three predictive variables that additionally incorporate interest rate proxies. We discuss below the results, but we do not report the details for brevity.

testing methods. We exploit the properties of our robust testing method to identify such data points. Figure 12 plots the time series of Huber weights estimated by the robust estimator (15) of the predictive regression model:

$$h(Z_t) := \min \left(1, \frac{c}{\|(Y_t - X'_{t-1} \hat{\alpha}_n^R) X_{t-1}\|} \right); \quad (22)$$

where $Z_t = (\ln(R_t); X_{t-1})'$, $X_{t-1} = (1; \ln(D_{t-1}=P_{t-1}))'$, and $\hat{\alpha}_n^R$ is the robust Huber estimator of parameter vector $\alpha = (\alpha_0; \alpha_1)'$. By definition, the data-driven weight $0 \leq h(Z_t) \leq 1$ is designed to reduce the influence of potential anomalous observations on the estimation results. Weights clearly below one indicate an anomalous data-point, while weights equal to one indicate unproblematic observations for the postulated model. Therefore, the value of weight (22) provides a standard way for highlighting potential anomalous observations that might be excessively influential for the fit of the predictive regression model (21); see, e.g., Hampel, Ronchetti, Rousseeuw and Stahel (1986).

We find that subperiod 1998-2002 is characterized by a cluster of infrequent anomalous observations, which are likely related to the abnormal stock market performance during the NASDAQ bubble in the second half of the 1990s. Similarly, we find a second cluster of anomalous observations in subperiod 2008-2010, which is linked to the extraordinary events of the recent financial crisis. Overall, anomalous observations are less than 4.2% of the whole data sample and they explain the failure of conventional testing methods in uncovering hidden predictability structures in these sample periods.

We find that the most influential observation before 1995 is November 1987, following the Black Monday on October 19 1987. During the subperiod 1998-2002, the most influential observation is October 2001, reflecting the impact on financial markets of the terrorist attack on September 11 2001. Finally, the most anomalous observation in the whole sample period 1980-2010 is October 2008, following the Lehman Brothers default on September 15 2008. The impact on the test results of the Lehman Brothers default emerges also in Figure 7, where nonrobust resampling methods no longer reject \mathcal{H}_0 in 2009. In contrast, robust tests still find

significance evidence in favor of predictability.

5.2 Two-Predictor Model

We extend our empirical study to two-predictor regression models. This approach has several purposes. First, it can serve to assess the incremental predictive ability of the dividend yield, in relation to other well-known competing predictive variables. Second, it is useful to verify the power properties of robust bootstrap and subsampling tests in settings with several predictive variables.

Section 5.2.1 borrows from Bollerslev, Tauchen and Zhou (2009) and studies the joint predictive ability of the dividend yield and the variance risk premium. Section 5.2.2 follows the two-predictor model in Santos and Veronesi (2006), which considers the ratio of labor income to consumption as an additional predictive variable to the dividend yield.

5.2.1 Bollerslev, Tauchen and Zhou

We consider again monthly S&P 500 index and dividend data between January 1990 and December 2010, and test the predictive regression model:

$$\frac{1}{k} \ln(R_{t+k,t}) = \alpha_0 + \alpha_1 \ln\left(\frac{D_t}{P_t}\right) + \alpha_2 VRP_t + \varepsilon_{t+k,t}, \quad (23)$$

where $\ln(R_{t+k,t}) := \ln(R_{t+1}) + \dots + \ln(R_{t+k})$ and the variance risk premium $VRP_t := IV_t - RV_t$ is defined by the difference of the S&P 500 index option-implied volatility at time t , for one month maturity options, and the ex-post realized return variation over the period $[t-1; t]$. Bollerslev, Tauchen and Zhou (2009) show that the variance risk premium is the most significant predictive variable of market returns over a quarterly horizon. Therefore, we test the predictive regression model (23) for $k = 4$.

Let α_{01} and α_{02} denote the true values of parameters α_1 and α_2 , respectively. Using the conventional bootstrap and subsampling tests, as well as our robust bootstrap and subsampling tests, we first test the null hypothesis of no return predictability by the dividend yield, \mathcal{H}_{01} :

$\alpha_1 = 0$.

Figure 8 plots the 90%-confidence intervals for parameter α_1 , based on rolling windows of 180 monthly observations in sample period 1990-2010, while Table 4 collects the detailed point estimates and testing results. We find again that the robust tests always clearly reject the null of no predictability at the 5%-significance level. In contrast, the conventional bootstrap and subsampling tests produce weaker and more ambiguous results, with uniformly lower p -values (larger confidence intervals) and a non-rejection of the null of no predictability at the 5%-level in period 1994-2009. Since the Bonferroni approach in Campbell and Yogo (2006) is defined for single-predictor models, we cannot apply this method in model (23). Unreported results for the multi-predictor testing method in Amihud, Hurvich and Wang (2008) show that for data windows following window 1993-2008 also the bias-corrected method cannot reject null hypothesis \mathcal{H}_{01} at the 10% significance level.

By inspecting the Huber weights (22), implied by the robust estimation of the predictive regression model (23), we find again a cluster of infrequent anomalous observations, both during the NASDAQ bubble and the recent financial crisis. Also in this setting, the most influential observation is October 2008, reflecting the Lehman Brothers default on September 15 2008. The impact of these anomalous observations emerges also in Figure 8, explaining the large estimated confidence intervals of nonrobust tests and their non-rejection of \mathcal{H}_{01} in subperiod 1994-2009.

Finally, we test the hypothesis of no predictability by the variance risk premium, $\mathcal{H}_{02} : \alpha_2 = 0$. Figure 9 plots the resulting confidence intervals for parameter α_2 , while Table 4 reports the estimates and testing results. In contrast to the previous evidence, we find that all tests under investigation clearly reject \mathcal{H}_{02} at the 5%-significance level, thus confirming the remarkable return forecasting ability of the variance risk premium noticed in Bollerslev, Tauchen and Zhou (2009), as well as the international evidence reported in Bollerslev, Marrone, Xu and Zhou (2012).

5.2.2 Santos and Veronesi

We finally focus on the two-predictor regression model proposed in Santos and Veronesi (2006):

$$\ln(R_t) = \alpha_0 + \alpha_1 \ln\left(\frac{D_{t-1}}{P_{t-1}}\right) + \alpha_2 S_{t-1} + \epsilon_t \quad (24)$$

where $S_{t-1} = w_{t-1} = C_{t-1}$ is the share of labor income to consumption. We make use of quarterly returns on the value weighted CRSP index, which includes NYSE, AMEX, and NASDAQ stocks, in the sample period Q1,1955-Q4,2010. The dividend time-series is also obtained from CRSP, while the risk free rate is the three-months Treasury bill rate. Labor income and consumption are obtained from the Bureau of Economic Analysis.²³

Let α_{01} and α_{02} denote the true values of parameters α_1 and α_2 , respectively. Using bootstrap and subsampling tests, as well as our robust testing method, we first test the null hypothesis of no predictability by the dividend yield, $\mathcal{H}_{01} : \alpha_{01} = 0$. Figure 10 plots the 90%-confidence intervals for parameter α_{01} based on rolling windows of 180 quarterly observations in sample period 1950-2010, while Table 5 collects detailed point estimates and test results for the four subperiods 1950-1995, 1955-2000, 1960-2005, 1965-2010. We find again that our robust tests always clearly reject \mathcal{H}_{01} at the 5%-significance level. In contrast, conventional tests produce more ambiguous results and cannot reject at the 10%-significance level the null hypothesis \mathcal{H}_{01} for subperiod 1955-2000.

Figure 11 reports the 90%-confidence intervals estimated in tests of the null hypothesis of no predictability by the labor income proxy, $\mathcal{H}_{02} : \alpha_{02} = 0$. Table 5 summarizes estimation and testing results. While the conventional tests produce a weak and mixed evidence of return predictability using labor income proxies, e.g., by not rejecting \mathcal{H}_{02} at the 10%-level in subperiod 1950-1995, the robust tests produce once more a clear and consistent predictability evidence for all sample periods.

The clusters of anomalous observations (less than 4.6% of the data in the full sample),

²³As in Lettau and Ludvigson (2001), labor income is defined as wages and salaries, plus transfer payments, plus other labor income, minus personal contributions for social insurance, minus taxes. Consumption is defined as non-durables plus services.

highlighted by the estimated weights in Figure 14, again indicate that conventional tests might fail to uncover hidden predictability structures using samples of data that include observations from the NASDAQ bubble or the recent financial crisis, a feature that have already noted also in Santos and Veronesi (2006) and Lettau and Van Nieuwerburgh (2007) from a different angle. In such contexts, the robust bootstrap and subsampling tests are again found to control well the potential damaging effects of anomalous observations, by providing a way to consistently uncover hidden predictability features also when the data may only approximately follow the given predictive regression model.

6 Conclusion

A large literature studies the predictive ability of a variety of economic variables for future market returns. Several useful testing approaches for testing the null of no predictability in predictive regressions with correlated errors and nearly integrated regressors have been proposed, including tests that rely on nonparametric Monte Carlo simulation methods, such as the bootstrap and the subsampling. All these methods improve on the conventional asymptotic tests under the ideal assumption of an exact predictive regression model. However, we find by Monte Carlo evidence that even small violations of such assumptions, generated by a small fraction of anomalous observations, can result in large deteriorations in the reliability of all these tests.

To systematically understand the problem, we characterize theoretically the robustness properties of bootstrap and subsampling tests of predictability in a time series context, using the concept of quantile breakdown point, which is a measure of the global resistance of a testing procedure to outliers. We obtain general quantile breakdown point formulas, which highlight an important non-resistance of these tests to anomalous observations that might infrequently contaminate the predictive regression model, thus confirming the fragility detected in our Monte Carlo study.

We propose a more robust testing method for predictive regressions with correlated errors and nearly integrated regressors, by introducing a novel general class of fast and robust

bootstrap and subsampling procedures for times series, which are applicable to linear and non-linear predictive regression models at sustainable computational costs. The new bootstrap and subsampling tests are resistant to anomalous observations in the data and imply more robust confidence intervals and inference results. We demonstrate by Monte Carlo simulations their good resistance to outliers and their improved finite-sample properties in presence of anomalous observations.

In our empirical study for US stock market data, we study single-predictor and multi-predictor models, using well-known predictive variables in the literature, such as the market dividend yield, the difference between index option-implied volatility and realized volatility (Bollerslev, Tauchen and Zhou, 2009) and the share of labor income to consumption (Santos and Veronesi, 2006).

First, using the robust tests we find clear-cut evidence that the price-dividend ratio is a robust predictive variable for market returns, in each subperiod and for each sampling frequency and forecasting horizon considered. In contrast, tests based on bias-corrections, near-to-unity asymptotics or standard resampling procedures provide more ambiguous findings, by not rejecting the null of no predictability in a number of cases.

Second, we find that the difference between option-implied volatility and realized volatility is a robust predictive variable of future market returns at quarterly forecasting horizons, both using robust and nonrobust testing methods. This finding confirms the remarkable return forecasting ability of the variance risk premium, first noticed in Bollerslev, Tauchen and Zhou (2009).

Third, we find that conventional testing approaches deliver an ambiguous evidence of return predictability by proxies of labor income, which is either absent or weak in the sample periods 1955-2000 and 1965-2010, respectively. In contrast, the null of no predictability is always clearly rejected using the robust testing approach, indicating that the weak findings of the conventional tests are likely deriving from their low ability to detect predictability structures in presence of small sets of anomalous observations.

Fourth, we exploit the properties of our robust tests to identify potential anomalous observations that might explain the diverging conclusions of robust and nonrobust methods. We

nd a fraction of less than about 5% of anomalous observations in the data, which tend to cluster during the NASDAQ bubble and the more recent financial crisis. Anomalous data points, including the Lehman Brothers default on September 2008, the terrorist attack of September 2001, the Black Monday on October 1987 and the Dot-Com bubble collapse in August 2002, are responsible for the failure of conventional testing methods in uncovering the hidden predictability structures for these sample periods.

Finally, while our bootstrap and subsampling tests have been developed in the context of standard predictive systems with autocorrelated regressors, our approach is extendable also to more general settings, including potential nonlinear predictive relations or unobserved state variables. For instance, van Binsbergen and Koijen (2010) propose a latent-variable approach and a Kalman filter to estimate a present value model with hidden and persistent expected return and dividend growth, in order to formulate powerful tests for the joint predictability of stock returns and dividend growth. The application of our robust bootstrap and subsampling tests in the context of such present value models is an interesting avenue for future research.

Appendix A: Robust Bootstrap and Subsampling Tests of Predictability

We present in more details how to compute the robust resampling distribution (17) and construct robust confidence intervals for the components of a general d -dimensional parameter θ , where $d \geq 1$. The extension to the nonstudentized distributions (16) is straightforward. We construct symmetric resampling confidence intervals for the parameter of interest.²⁴

Let $\theta = (\beta; \gamma)'$ and let $z_{(n)} = (z_1; \dots; z_n)$ be an observation sample generated according to the multi-predictor regression model

$$\begin{aligned} y_t &= \beta + \gamma' x_{t-1} + u_t; \\ x_t &= R x_{t-1} + v_t; \end{aligned}$$

where $z_t = (y_t; w_{t-1}')'$, $w_{t-1} = (1; x_{t-1}')'$, β is a d -dimensional parameter vector and R is a $d \times d$ parameter matrix. First, we compute the robust Huber estimator $\hat{\theta}_n^R = (\hat{\beta}_n^R, \hat{\gamma}_n^R)'$ as the solution of $\psi_{n,c}(z_{(n)}; \hat{\theta}_n^R) = 0$, where the estimating function $\psi_{n,c}$ is defined in (14) and $c > 0$ is a tuning constant selected through the data-driven method introduced in Appendix B below. Note that the asymptotic variance of the robust M-estimator $\hat{\theta}_n^R$ is given by

$$V^R = J'^R V^R J^R, \quad (25)$$

where $J^R = (\lim_{n \rightarrow \infty} E [\nabla_{\theta} \psi_{n,c}(z_{(n)}; \theta_0)])^{-1}$, $V^R = \lim_{n \rightarrow \infty} \text{Var} \left[\frac{1}{\sqrt{n}} \sum_{i=1}^n g_c(z_{(i)}; \theta_0) \right]$, and $\nabla_{\theta} \psi_{n,c}(z_{(n)}; \theta_0)$ denotes the derivative of function $\psi_{n,c}$ with respect to parameter θ with $\theta_0 =$

²⁴Hall (1988) and more recent contributions, as for instance Politis, Romano and Wolf (1999), highlight a better accuracy of symmetric confidence intervals, which even in asymmetric settings can be shorter than asymmetric confidence intervals. Mikusheva (2007) and Andrews and Guggenberger (2010) also show that because of a lack of uniformity in pointwise asymptotics, nonsymmetric subsampling confidence intervals for autoregressive models can imply a distorted asymptotic size, which is instead correct for symmetric confidence intervals.

$(\mathbf{0}, \mathbf{0})'$. We compute the estimator $\hat{\mathbf{V}}_n^R = \hat{\mathbf{J}}_n^R \hat{\mathbf{V}}_n^R \hat{\mathbf{J}}_n^R$ of the asymptotic variance \mathbf{V}^R where

$$\hat{\mathbf{J}}_n^R = \left[\nabla_{\theta} g_{n,c}(Z_{(n)}; \hat{\theta}_n^R) \right]^{-1}; \quad (26)$$

$$\hat{\mathbf{V}}_n^R = \frac{1}{n} \sum_{i=1}^n g_c(Z_{i, \cdot}; \hat{\theta}_n^R) g_c(Z_{i, \cdot}; \hat{\theta}_n^R)'; \quad (27)$$

respectively.²⁵ Let $(p)^{(j)}$ denotes the j -th component of a d -dimensional vector p , $1 \leq j \leq d$. To construct symmetric confidence intervals for each j -th component of the parameter θ , we compute an estimator of the sampling distribution of $(T_{n,|\cdot|})^{(j+1)} := \frac{\sqrt{n}}{(\hat{\theta}_n^R)^{(j+1)}} \left| \left(\hat{\theta}_n^R - \mathbf{0} \right)^{(j+1)} \right|$, by applying our robust resampling approach, where the subscript $|\cdot|$ indicates that we consider the absolute value of each component of the statistic T_n , and $(\hat{\theta}_n^R)^{(j+1)}$ is the square root of the $(j+1)$ -th diagonal component of matrix $\hat{\mathbf{V}}_n^R$.

More precisely, let $Z_{(n,m)}^{K*}$ be a block bootstrap ($K = B$) or subsampling ($K = S$) random sample based on blocks of size m , where m is selected through the data-driven method introduced in Appendix B below. For each component $1 \leq j \leq d$, we estimate the sampling distribution of $(T_{n,|\cdot|})^{(j+1)}$ through the robust resampling distribution

$$(L_{n,m,|\cdot|}^{K*}(x))^{(j+1)} = \frac{1}{N} \sum_{s=1}^N \mathbb{I} \left(\frac{\sqrt{k}}{(\hat{\mathbf{V}}_{k,s}^{R*})^{(j+1)}} \left| \left([\nabla_{\theta} g_{k,c}(Z_{(n,m),s}^{K*}; \hat{\theta}_n^R)]^{-1} g_{k,c}(Z_{(n,m),s}^{K*}; \hat{\theta}_n^R) \right)^{(j+1)} \right| \leq x \right); \quad (28)$$

where $k = n$ for the block bootstrap ($K = B$), while $k = m$ for the subsampling ($K = S$), $(\hat{\mathbf{V}}_{k,s}^{R*})^{(j+1)}$ is the square root of the $(j+1)$ -th diagonal component of matrix $\hat{\mathbf{V}}_{k,s}^{R*}$, and $\hat{\mathbf{V}}_{k,s}^{R*} = \hat{\mathbf{J}}_{k,s}^{R*} \hat{\mathbf{V}}_{k,s}^{R*} \hat{\mathbf{J}}_{k,s}^{R*}$ with

$$\hat{\mathbf{J}}_{k,s}^{R*} = \left[\nabla_{\theta} g_{k,c}(Z_{(n,m),s}^{K*}; \hat{\theta}_n^R) \right]^{-1}; \quad (29)$$

$$\hat{\mathbf{V}}_{k,s}^{R*} = \frac{1}{k} \sum_{i=1}^k g_c(Z_{i,s}; \hat{\theta}_n^R) g_c(Z_{i,s}; \hat{\theta}_n^R)'. \quad (30)$$

²⁵For the sake of brevity, we assume that $E[g_c(z_i, \theta_0)g_c(z_j, \theta_0)'] = 0$, $i \neq j$, and consequently we consider $\hat{\mathbf{V}}_n^R$ as estimator of \mathbf{V}^R . However, if this assumption is not satisfied, then $\hat{\mathbf{V}}_n^R$ has to be replaced with the Newey-West covariance estimator.

Finally, let $t \in (0; 1)$, and let $(Q_{t,n,m}^{K*})^{(j+1)}$ be the t -quantile of the block bootstrap or subsampling empirical distribution (28), $K = B; S$, respectively. Then, the symmetric t -confidence interval for the j -th component $(\beta_0)^{(j)}$ of β_0 is given by

$$(CI_t)^{(j)} = \left[\binom{\hat{\beta}_n^R}{n}^{(j+1)} - \binom{\hat{\beta}_n^R}{n}^{(j+1)} (Q_{t,n,m}^{K*})^{(j+1)}, \binom{\hat{\beta}_n^R}{n}^{(j+1)} + \binom{\hat{\beta}_n^R}{n}^{(j+1)} (Q_{t,n,m}^{K*})^{(j+1)} \right]; \quad (31)$$

We summarize our robust approach in the following steps.

- (1) Compute $\hat{\beta}_n^R = (\hat{\beta}_{ROB}^R; \hat{\beta}_{ROB}^R)'$, as the solution of (14), where c is selected using the data-driven method introduced in Appendix B below.
- (2) Compute $\hat{\beta}_n^R = \hat{J}_n^R \hat{V}_n^R \hat{J}_n^R$, where \hat{J}_n^R and \hat{V}_n^R are defined in (26) and (27), respectively.
- (3B) For the robust block bootstrap, generate $B_B = 999$ random bootstrap samples based on the overlapping blocks $(Z_i; \dots; Z_{i+m-1})$, $i = 1; \dots; n-m+1$, where m is selected according to the data-driven method introduced in Appendix B below.
- (3S) For the robust subsampling, consider the $B_S = n-m+1$ overlapping blocks $(Z_i; \dots; Z_{i+m-1})$, $i = 1; \dots; n-m+1$, where m is selected according to the data-driven method introduced in Appendix B below.
- (4) Compute the robust resampling distribution (28).
- (5) The robust symmetric t -confidence interval for the j -th component $(\beta_0)^{(j)}$ of β_0 is given by (31).

Appendix B: Data-Driven Selection of the Block Size and the Robust Estimating Function Bound c

The implementation of our robust resampling methods requires the selection of the block size m and the degree of robustness c of the estimating function (14). To this end, we propose a data-driven procedure for the choice of the block size m and the estimating function bound c , by extending the calibration method (CM) discussed in Romano and Wolf (2001) in relation to subsampling procedures.

Let $\mathcal{MC} := \{(m; c) | m \in \mathcal{M}; c \in \mathcal{C}\}$, where $\mathcal{M} := \{m_{\min} < \dots < m_{\max}\}$ and $\mathcal{C} := \{c_{\min} < \dots < c_{\max}\}$ are the sets of admissible block sizes and estimating functions bounds, respectively. Let $T_n^{NS} = \sqrt{n} \left(\hat{\tau}_n^R - \tau_0 \right)$ be the nonstudentized statistic of interest, where $\hat{\tau}_n^R$ is the robust Huber estimator solution of Equation (14) with $c = c_1$ fixed, as preliminary value of the estimating function bound. Furthermore, let $(X_1^*:::; X_n^*)$ be a block bootstrap sample generated from $(X_1:::; X_n)$, with the block size $m \in \mathcal{M}$. For each bootstrap sample, compute a t -subsampling (or bootstrap) confidence interval $CI_{t,(m,c)}$ as described in Appendix A according to block size $m \in \mathcal{M}$ and bound $c \in \mathcal{C}$. The data-driven block size and estimating function bound according to the calibration method are defined as

$$(m; c)_{CM} := \arg \inf_{(m,c) \in \mathcal{MC}} \left\{ \left| t - P^* \left[\hat{\tau}_n^R \in CI_{t,(m,c)} \right] \right| \right\}; \quad (32)$$

where, by definition, $\arg \inf(\emptyset) := \infty$, and P^* denotes the probability with respect to the bootstrap distribution. By definition, $(m; c)_{CM}$ is the pair for which the bootstrap probability of the event $\{ \hat{\tau}_n^R \in CI_{t,(m,c)} \}$ is as near as possible to the nominal level t of the confidence interval. The extension to the studentized statistic $T_n = \sqrt{n} \left([\hat{\tau}_n^R]^{-1/2} (\hat{\tau}_n^R - \tau_0) \right)$ is straightforward.

We summarize the calibration method for the selection of the block size m and estimating function bound c in the following steps.

- (1) Compute $\hat{\tau}_n^R = (\hat{\tau}_n^{R,1}, \hat{\tau}_n^{R,2})'$, as the solution of (14), with $c = c_1$ as preliminary value of the estimating function bound.

- (2) For each $m \in \mathcal{M}$, generate K random bootstrap samples $(z_1^*; \dots; z_n^*)$ based on overlapping blocks of size m .
- (3) For each random bootstrap sample $(z_1^*; \dots; z_n^*)$ and $c \in \mathcal{C}$, compute confidence intervals $CI_{t,(m,c)}$ for the parameter $\hat{\theta}_n^R$, by applying steps (1)-(5) of the algorithm in the previous Appendix A.
- (4) For each pair $(m; c) \in \mathcal{MC}$ compute $h(m; c) = J\{\hat{\theta}_n^R \in CI_{t,(m,c)}\} = K$.
- (5) The data-driven block size and estimating function bound according to the calibration method are defined as

$$(m; c)_{CM} := \arg \inf_{(m,c) \in \mathcal{MC}} \{|t - h(m; c)|\} :$$

Appendix C: Proofs

Proof of Theorem 1. We first consider the subsampling and focus on formula (7). The value $\frac{\lceil mb \rceil}{n}$ is the smallest fraction of outliers, that causes the breakdown of statistic T in a block of size m . Therefore, the first inequality is satisfied.

For the second inequality of formula (7), we denote by $X_{(m),i}^N = (X_{(i-1)m+1} : \dots : X_{im})$, $i = 1, \dots, r$ and $X_{(m),i}^O = (X_i : \dots : X_{i+m-1})$, $i = 1, \dots, n-m+1$, the nonoverlapping and overlapping blocks of size m , respectively. Given the original sample $X_{(n)}$, for the first nonoverlapping block $X_{(m),1}^N$, consider the following type of contamination:

$$X_{(m),1}^N = (X_1 : \dots : X_{m-\lceil mb \rceil}; Z_{m-\lceil mb \rceil+1} : \dots : Z_m); \quad (33)$$

where X_i , $i = 1, \dots, m - \lceil mb \rceil$ and Z_j , $j = m - \lceil mb \rceil + 1, \dots, m$, denote the noncontaminated and contaminated points, respectively. By construction, the first $m - \lceil mb \rceil + 1$ overlapping blocks $X_{(m),i}^O$, $i = 1, \dots, m - \lceil mb \rceil + 1$, contain $\lceil mb \rceil$ outliers. Consequently, $T(X_{(m),i}^O) = +\infty$, $i = 1, \dots, m - \lceil mb \rceil + 1$. Assume that the first $p < r - 1$ nonoverlapping blocks $X_{(m),i}^N$, $i = 1, \dots, p$, have the same contamination as in (33). Because of this contamination, the number of statistics $T_{n,m}^{OS*}$ which diverge to infinity is $mp - \lceil mb \rceil + 1$.

$Q_{t,n,m}^{OS*} = +\infty$, when the proportion of statistics $T_{n,m}^{OS*}$ with $T_{n,m}^{OS*} = +\infty$ is larger than $(1 - t)$. Therefore,

$$b_{t,n,m}^{OS} \leq \inf_{\{p \in \mathbb{N}, p \leq r-1\}} \left\{ p \cdot \frac{\lceil mb \rceil}{n} \left| \frac{mp - \lceil mb \rceil + 1}{n - m + 1} > 1 - t \right. \right\}.$$

Finally, we consider formula (8). The proof of the first inequality in formula (8) follows the same lines as the proof of the first inequality in the formula (7). We focus on the second inequality.

Consider $X_{(m),i}^N$, $i = 1, \dots, r$. Assume that p_2 of these nonoverlapping blocks are contaminated with exactly p_1 outliers for each block, while the remaining $(r - p_2)$ are noncontaminated

(0 outlier), where $p_1, p_2 \in \mathbb{N}$ and $p_1 \leq m, p_2 \leq r - 1$. Moreover, also assume that the contamination of the p_2 contaminated blocks has the structure defined in (33). The block bootstrap constructs a n -sample randomly selecting with replacement r overlapping blocks of size m . Let X be the random variable which denotes the number of contaminated blocks in the random bootstrap sample. It follows that $X \sim \text{BIN}(r; \frac{mp_2 - p_1 + 1}{n - m + 1})$.

By Equation (5), $Q_{t,n,m}^{OB*} = +\infty$, when the proportion of statistics $T_{n,m}^{OB*}$ with $T_{n,m}^{OB*} = +\infty$ is larger than $(1 - t)$. The smallest number of outliers such that $T_{n,m}^{OB*} = +\infty$ is by definition nb . Let $p_1, p_2 \in \mathbb{N}; p_1 \leq m; p_2 \leq r - 1$. Consequently,

$$b_{t,n,m}^{OB} \leq \frac{1}{n} \cdot \left[\inf_{\{p_1, p_2\}} \left\{ p = p_1 \cdot p_2 \left| P \left(\text{BIN} \left(r; \frac{mp_2 - p_1 + 1}{n - m + 1} \right) \geq \left\lceil \frac{nb}{p_1} \right\rceil \right) > 1 - t \right\} \right]:$$

This concludes the proof of Theorem 1. ■

Proof of Corollary 3. Consider the robust fast approximation of $\hat{\theta}_{n,m}^{K*} - \hat{\theta}_n$ given by:

$$-[\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)]^{-1} \cdot k(X_{(n,m),s}^{K*}; \hat{\theta}_n); \quad (34)$$

where $k = n$ or $k = m$, $K = B; S$. Assuming a bounded estimating function, Expression (34) may degenerate only when either (i) $\hat{\theta}_n \in \mathbb{R}$ or (ii) matrix $[\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)]$ is singular, i.e., $\det([\nabla_{\theta} \psi_n(X_{(n)}; \hat{\theta}_n)]) = 0$. If (i) and (ii) are not satisfied, then, quantile $Q_{t,n,m}^{K*}$ is bounded, for all $t \in (0; 1)$. Let b be the breakdown point of $\hat{\theta}_n$ and $b_{\nabla\psi}$ the smallest fraction of outliers in the original sample such that condition (ii) is satisfied. Then, the breakdown point of $Q_{t,n,m}^{K*}$ is $b_{t,n,m}^K = \min(b; b_{\nabla\psi})$. ■

Proof of Corollary 4. First note that the breakdown point of the robust estimator $\hat{\theta}_n^R$ defined in (14) is $b = 0.5$; see, e.g., Huber (1984). Therefore, we have only to focus on the breakdown point of matrix $\nabla_{\theta}(\psi_{n,c}(Z_{(n)}; \hat{\theta}_n^R))$. Without loss of generality, assume that $\theta = (\beta; \gamma) \in \mathbb{R}^2$. Consider the function

$$g_c(y_t; w_{t-1}; \gamma) = (y_t - \gamma' w_{t-1}) w_{t-1} \cdot \min \left(1; \frac{c}{\| (y_t - \gamma' w_{t-1}) w_{t-1} \|} \right); \quad (35)$$

Using some algebra, we can show that

$$\nabla_{\theta} g_c(y_t; w_{t-1}) = \begin{cases} -(1; x_{t-1})'(1; x_{t-1}); & \text{if } \|(y_t - \hat{w}_{t-1})w_{t-1}\| \leq c; \\ \mathbb{O}_{2 \times 2}; & \text{if } \|(y_t - \hat{w}_{t-1})w_{t-1}\| > c; \end{cases} \quad (36)$$

where $\mathbb{O}_{2 \times 2}$ denotes the 2×2 null matrix. It turns out that by construction the matrix $\nabla_{\theta}(\hat{z}_{n,c}(z_{(n)}; \hat{n}^R))$ is semi-positive definite, and in particular $\det(\nabla_{\theta}(\hat{z}_{n,c}(z_{(n)}; \hat{n}^R))) = 0$, only when $\|(y_t - \hat{w}_n^R w_{t-1})w_{t-1}\| > c$, for all the observations $(y_t; w_{t-1})'$, i.e., $b_{\nabla\psi_c} = 1$. Therefore, using Corollary 3 we obtain that $b_{t,n,m}^K = \min(0.5; 1) = 0.5$, $K = B; S$. ■

Proof of Theorem 5. Consider the resampling distribution (17). Let $z_{(n,m),s}^{K*} = (z_1^* \cdots z_k^*)$ denote a random bootstrap ($K = B$ and $k = n$) or subsampling ($K = S$ and $k = m$) sample. Since the estimating function $\hat{z}_{n,c}$ is bounded, it turns out that

$$T_{n,m,s}^{K*} := [\hat{z}_{k,s}^{R*}]^{-1/2} [\nabla_{\theta} \hat{z}_{k,c}(z_{(n,m),s}^{K*}; \hat{n}^R)]^{-1} \hat{z}_{k,c}(z_{(n,m),s}^{K*}; \hat{n}^R); \quad (37)$$

may degenerate when (i) $\det(\hat{z}_{k,s}^{R*}) = 0$ or (ii) $\det(\nabla_{\theta} \hat{z}_{k,c}(z_{(n,m),s}^{K*}; \hat{n}^R)) = 0$. Moreover, also note that $\hat{z}_{k,s}^{R*} = \hat{J}_{k,s}^{R*} \hat{J}_{k,s}^{R*}$, where $\hat{J}_{k,s}^{R*} = [\nabla_{\theta} \hat{z}_{k,c}(z_{(n,m),s}^{K*}; \hat{n}^R)]^{-1}$. Because of Equations (29), (30), (36), it turns out that cases (i) and (ii) can be satisfied only when $\|(y_t^* - \hat{w}_n^R w_{t-1}^*)w_{t-1}^*\| > c$ for all random observations $z_i^* = (y_i^*; w_{i-1}^*)'$, $i = 1; \dots; k$, where $k = n; m$, for the bootstrap and subsampling, respectively.

For the original sample, consider following type of contamination

$$z_{(n)} = (z_1; \dots; z_j; C_{j+1}; \dots; C_{j+p}; z_{j+p+1}; \dots; z_n); \quad (38)$$

where z_i , $i = 1; \dots; j$ and $i = j + p + 1; \dots; n$ and C_i , $i = j + 1; \dots; j + p$, denote the noncontaminated and contaminated points, respectively, where $p \geq m$. It turns out that all the $p - m + 1$ overlapping blocks of size m

$$(C_{j+i}; \dots; C_{j+i+m-1}); \quad (39)$$

$i = 1; \dots; p - m + 1$ contain only outliers. Therefore, for these $p - m + 1$ blocks we have that $\det(\nabla_{\theta} \mathcal{L}_{m,c}(C_{j+i}; \dots; C_{j+i+m-1}; \hat{\Lambda}_n^R)) = 0$, i.e., some components of vector (37) may degenerate to infinity. Moreover, $Q_{t,n,m}^{S*} = +\infty$ when the proportion of statistics $T_{n,m}^{S*}$ with $T_{n,m}^{S*} = +\infty$ is larger than $(1 - t)$. Therefore, $b_{t,n,m}^S = \inf_{\{p \in \mathbb{N}, m \leq p \leq n-m+1\}} \left\{ \frac{p}{n} \left| \frac{p-m+1}{n-m+1} > 1 - t \right. \right\}$, which proves the result in Equation (18).

For the result in Equation (19), note that because of the contamination defined in (38), by construction we have $p - m + 1$ overlapping blocks of size m with exactly m outliers, and $n - (p - m + 1)$ blocks with less than m outliers. Let X be the random variable which denotes the number of full contaminated blocks in the random bootstrap sample. It follows that $X \sim \text{BIN}\left(r; \frac{p-m+1}{n-m+1}\right)$. To imply (i) or (ii), all the random observations $(z_1^*; \dots; z_k^*)$ have to be outliers, i.e., $X = r$. By Equation (5), $Q_{t,n,m}^{B*} = +\infty$, when the proportion of statistics $T_{n,m}^{B*}$ with $T_{n,m}^{B*} = +\infty$ is larger than $(1 - t)$. Consequently,

$$b_{t,n,m}^B = \frac{1}{n} \cdot \left[\inf_{\{p \in \mathbb{N}, p \leq n-m+1\}} \left\{ p \left| P\left(\text{BIN}\left(r; \frac{p-m+1}{n-m+1}\right) = r\right) > 1 - t \right. \right\} \right]:$$

This concludes the proof. ■

References

- [1] Amihud, Y., and C.M. Hurvich, 2004. Predictive regressions: a reduced-bias estimation method. *Journal of Financial and Quantitative Analysis*, 39, 813{841.
- [2] Amihud, Y., Hurvich C.M., and Y. Wang, 2008. Multiple-predictor regressions: hypothesis testing. *The Review of Financial Studies*, 22, 413{434.
- [3] Amihud, Y., Hurvich C.M., and Y. Wang, 2010. Predictive regression with order-p autoregressive predictors. *Journal of Empirical Finance*, 17, 513{525.
- [4] Andrews, D., 2002. Higher-order improvements of a computationally attractive k-step bootstrap for extremum estimators. *Econometrica*, 70, 119{162.
- [5] Andrews, D., 2004. The block-block bootstrap: improved asymptotic refinements. *Econometrica*, 72, 673{700.
- [6] Andrews, D., and P. Guggenberger, 2010. Application of subsampling, hybrid and size-correction methods. *Journal of Econometrics*, 158, 285-305.
- [7] Ang, A., and G. Bekaert, 2007. Stock return predictability: is it there? *The Review of Financial Studies*, 20, 651-707.
- [8] Bollerslev, T., Marrone, J., Xu, L., and H. Zhou, 2012. Stock return predictability and variance risk premia: statistical inference and international evidence. Working paper.
- [9] Bollerslev, T., Tauchen, G., and H. Zhou, 2009. Expected stock returns and variance risk premia. *Review of Financial Studies*, 22, 4463{4492.
- [10] Campbell, J.Y., and R.J. Shiller, 1988. The dividend ratio model and small sample bias: a Monte Carlo study. *Economics Letters*, 29, 325{331.
- [11] Campbell, J.Y., and M. Yogo, 2006. Efficient tests of stock return predictability. *Journal of Financial Economics*, 81, 27{60.

- [12] Camponovo, L., Scaillet, O., and F. Trojani, 2012. Robust subsampling. *Journal of Econometrics*, 167, 197{210.
- [13] Carlstein, E., 1986. The use of subseries methods for estimating the variance of a general statistic from a stationary time series. *Annals of Statistics*, 14, 1171{1179.
- [14] Cavanagh, C., Elliott, G., and J.H. Stock, (1995). Inference in models with nearly integrated regressors. *Econometric Theory* 11, 1131{1147.
- [15] Choi, I., and T.K. Chue, 2007. Subsampling-based tests of stock-return predictability, Working paper.
- [16] Davidson, R., and J. McKinnon, 1999. Bootstrap testing in nonlinear models. *International Economic Review*, 40, 487{508.
- [17] Donoho, D. L. and P. J. Huber (1983) The notion of breakdown point, in *A Festschrift for Erich L. Lehmann* by Bickel, P. J., Doksum, K. A. and J. L. Hodges Jr. (eds.), 157-184, Wadsworth, Belmont, California.
- [18] Fama, E., and K. French, 1988. Dividend yields and expected stock returns. *Journal of Financial Economics*, 22, 3-25.
- [19] Gagliardini P., Trojani F., and G. Urga, (2005). Robust GMM tests for structural breaks. *Journal of Econometrics*, 129, 139-182.
- [20] Goetzmann, W.N., and P. Jorion, 1995. A longer look at dividend yields. *Journal of Business*, 68, No.4.
- [21] Genton, M., and A. Lucas, 2003. Comprehensive definitions of breakdown point for independent and dependent observations. *Journal of the Royal Statistical Society, Series B*, 65, 81{94.
- [22] Goncalves, S., and H. White, 2004. Maximum likelihood and the bootstrap for nonlinear dynamic models. *Journal of Econometrics*, 119, 199{220.

- [23] Goyal, A., and I. Welch, 2003. Predicting the equity premium with dividend ratios. *Management Science*, 49, 639{654.
- [24] Hall, P., 1985. Resampling a coverage process. *Stochastic Processes and their Applications*, 19, 259{269.
- [25] Hall, P., 1988. On symmetric bootstrap confidence intervals. *Journal of the Royal Statistical Society, Ser.B*, 50, 35{45.
- [26] Hall, P., and J. Horowitz, 1996. Bootstrap critical values for tests based on Generalized-Method-of-Moment estimators. *Econometrica*, 64, 891{916.
- [27] Hall, P., Horowitz J., and B.-Y. Jing, 1995. On blocking rules for the bootstrap with dependent data. *Biometrika*, 82, 561{574.
- [28] Hampel, F. R., (1971) A General Qualitative Definition of Robustness, *Annals of Mathematical Statistics*, 42, 1887-1896.
- [29] Hampel, F. R., Ronchetti, E. M., Rousseeuw, P. J., and W. A. Stahel, 1986. *Robust statistics. The approach based on influence functions*. Wiley, New York.
- [30] Heritier, S., and E. Ronchetti, 1994. Robust bounded-influence tests in general parametric model. *Journal of the American Statistical Association*, 89, 897-904.
- [31] Hong, H., and O. Scaillet, 2006. A fast subsampling method for nonlinear dynamic models. *Journal of Econometrics*, 133, 557-578.
- [32] Hu, F., and J. Kalbfleisch, 2000. The estimating function bootstrap. *Canadian Journal of Statistics*, 28, 449{499
- [33] Huber, P. J., 1981. *Robust Statistics*. Wiley, New York.
- [34] Huber, P.J., 1984. Finite sample breakdown point of M- and P-estimators. *Annals of Statistics*, 12, 119{126.

- [35] Kothari, S.P., and J. Shanken, 1997. Book-to-market, dividend yield, and expected market returns: a time series analysis. *Journal of Financial Economics*, 2, 169-203.
- [36] Kunsch, H., 1989. The jackknife and the bootstrap for general stationary observations. *Annals of Statistics*, 17, 1217{1241.
- [37] Jansson, M., and M.J. Moreira, 2006. Optimal inference in regression models with nearly integrated regressors. *Econometrica*, 74, 681{714.
- [38] Lettau, M., and S. Ludvigson, 2001. Consumption, aggregate wealth, and expected stock returns. *Journal of Finance*, 3, 815{849.
- [39] Lettau, M., and S. Van Nieuwerburgh, 2007. Reconciling the return predictability evidence: in-sample forecasts, out-of-sample forecasts, and parameter instability. *Review of Financial Studies*, 21, 1607{1652.
- [40] Lewellen, J., 2004. Predicting returns with financial ratios. *Journal of Financial Economics*, 74, 209{235.
- [41] Lucas, A., 1995. Unit root tests based on M-estimators. *Econometric Theory*, 11, 331{346.
- [42] Lucas, A., 1997. Cointegration testing using pseudolikelihood ratio tests. *Econometric Theory*, 13, 149{169.
- [43] Mancini, L., Ronchetti, E., and F. Trojani, 2005. Optimal conditionally unbiased bounded-influence inference in dynamic location and scale models. *Journal of the American Statistical Association*, 100, 628-641.
- [44] Mankiv, N.G., and M.D. Shapiro, 1986. Do we reject too often? Small sample properties of tests of rational expectation models. *Economics Letters* 20, 139-145.
- [45] Mikusheva, A., 2007. Uniform inference in autoregressive models. *Econometrica*, 75, 1411{1452.

- [46] Nelson, C.R., and M.J. Kim, 1993. Predictable stock returns: the role of small sample bias. *Journal of Finance*, 48, No.2, 641-661.
- [47] Politis, D. N., and J. P. Romano, 1994. The stationary bootstrap. *Journal of the American Statistical Association*, 89, 1303{1313.
- [48] Politis, D.N., Romano J.P., and M. Wolf, 1999. *Subsampling*. Springer, New York.
- [49] Polk, C., S. Thompson, and T. Vuolteenaho, 2006. Cross-sectional forecast of the equity premium. *Journal of Financial Economics*, 81, 101-141.
- [50] Romano, J. P., and M. Wolf, 2001. Subsampling intervals in autoregressive models with linear time trend. *Econometrica*, 69, 1283-1314.
- [51] Ronchetti, E., and F. Trojani, 2001. Robust inference with GMM estimators, *Journal of Econometrics*, 101, 37{69.
- [52] Roze , M., 1984. Dividend yields are equity risk premium. *Journal of Portfolio Management*, 11, 68-75.
- [53] Salibian-Barrera, M., and R. Zamar, 2002. Bootstrapping robust estimates of regression. *Annals of Statistics*, 30, No. 2, 556{582.
- [54] Salibian-Barrera, M., Van Aelst, S., and G. Willems, 2006. Principal components analysis based on multivariate MM estimators with fast and robust bootstrap. *Journal of the American Statistical Association*, 101, 1198{1211.
- [55] Salibian-Barrera, M., Van Aelst, S., and G. Willems, 2007. Fast and robust bootstrap. *Statistical Methods and Applications*, 17, 41{71
- [56] Santos, T., and P. Veronesi, 2006. Labor income and predictable stock returns. *Review of Financial Studies*, 19, 1{44.
- [57] Shao, J., and D. Tu, 1995. *The jackknife and bootstrap*. Springer, New York.

- [58] Shiller, R.J., 2000. *Irrational Exuberance*. Princeton University Press, Princeton, NJ.
- [59] Singh, K., 1998. Breakdown theory for bootstrap quantiles. *Annals of Statistics*, 26, 1719{1732.
- [60] Stambaugh, R.F., 1986. Bias in regressions with lagged stochastic regressors. Graduate School of Business, University of Chicago, Working Paper No. 156.
- [61] Stambaugh, R.F., 1999. Predictive regressions. *Jornal of Financial Economics*, 54, 375-421.
- [62] Torous, W., R. Valkanov, and S. Yan, 2004. On predicting stock returns with nearly integrated explanatory variables. *Journal of Business*, 77, 937-966.
- [63] van Binsbergen, J.H., and R.S. Koijen, 2010. Predictive regressions: A present-value approach. *Journal of Finance*, 65, 1439{1471.
- [64] Wolf, M., 2000. Stock returns and dividend yields revisited: a new way to look at an old problem. *Journal of Business and Economic Statistics*, 18, 18-30.




Figure 1: **Power curves of bias-corrected and local-to-unity asymptotics.** We plot the proportion of rejections of the null hypothesis $\mathcal{H}_0 : \beta_0 = 0$, when the true parameter value is $\beta_0 \in [0, 0.15]$. In the left panel, we consider the bias-corrected method proposed in Amihud, Hurvich and Wang (2008), while in the right panel we consider the Bonferroni approach for the local-to-unity asymptotic theory introduced in Campbell and Yogo (2006). We consider non-contaminated samples (straight line) and contaminated samples (dashed line).




Figure 2: **Power curves of block bootstrap and subsampling.** We plot the proportion of rejections of the null hypothesis $\mathcal{H}_0 : \beta_0 = 0$, when the true parameter value is $\beta_0 \in [0, 0.15]$. In the left panel, we consider the block bootstrap, while in the right panel we consider the subsampling. We consider non-contaminated samples (straight line) and contaminated samples (dashed line).




Figure 3: **Power curves of robust block bootstrap and robust subsampling.** We plot the proportion of rejections of the null hypothesis $\mathcal{H}_0 : \beta_0 = 0$, when the true parameter value is $\beta_0 \in [0, 0.15]$. In the left panel, we consider our robust block bootstrap, while in the right panel we consider our robust subsampling. We consider non-contaminated samples (straight line) and contaminated samples (dashed line).




Figure 4: **Sensitivity analysis of bias-corrected and local-to-unity asymptotics.** We plot the percentage of increase of the confidence interval lengths with respect to variation of y_{max} , in each Monte Carlo sample, within the interval $[0, 5]$. In the left panel, we consider the bias-corrected method proposed in Amihud, Hurvich and Wang (2008), while in the right panel we consider the Bonferroni approach for the local-to-unity asymptotic theory introduced in Campbell and Yogo (2006).




Figure 5: **Sensitivity analysis of block bootstrap and subsampling.** We plot the percentage of increase of the confidence interval lengths with respect to variation of y_{max} , in each Monte Carlo sample, within the interval $[0, 5]$. In the left panel, we consider the block bootstrap, while in the right panel we consider the subsampling.




Figure 6: **Sensitivity analysis of robust block bootstrap and robust subsampling.** We plot the percentage of increase of the confidence interval lengths with respect to variation of y_{max} , in each Monte Carlo sample, within the interval $[0, 5]$. In the left panel, we consider our robust block bootstrap, while in the right panel we consider our robust subsampling.




Figure 7: **Upper and lower bounds of the confidence intervals.** We plot the upper and lower bound of the 90% confidence intervals for the parameter β in the predictive regression model (21). We consider rolling windows of 180 observations for the period 1980-2010. In the first line, we present the bias-corrected method (left panel) and the Bonferroni approach (right panel). In the second line, we consider the classic bootstrap (left panel) and the classic subsampling (right panel), while in the third line we consider our robust bootstrap (left panel) and our robust subsampling (right panel).




Figure 8: **Upper and lower bounds of the confidence intervals.** We plot the upper and lower bound of the 90% confidence intervals for the parameter β_1 in the predictive regression model (23). We consider rolling windows of 180 observations for the period 1990-2010. In the top line, we present the classic bootstrap (left panel) and the classic subsampling (right panel), while in the bottom line we consider our robust bootstrap (left panel) and our robust subsampling (right panel).




Figure 9: **Upper and lower bounds of the confidence intervals.** We plot the upper and lower bound of the 90% confidence intervals for the parameter β_2 in the predictive regression model (23). We consider rolling windows of 180 observations for the period 1990-2010. In the top line, we present the classic bootstrap (left panel) and the classic subsampling (right panel), while in the bottom line we consider our robust bootstrap (left panel) and our robust subsampling (right panel).




Figure 10: **Upper and lower bounds of the confidence intervals.** We plot the upper and lower bound of the 90% confidence intervals for the parameter β_1 in the predictive regression model (24). We consider rolling windows of 180 observations for the period 1950-2010. In the top line, we present the classic bootstrap (left panel) and the classic subsampling (right panel), while in the bottom line we consider our robust bootstrap (left panel) and our robust subsampling (right panel).




Figure 11: **Upper and lower bounds of the confidence intervals.** We plot the upper and lower bound of the 90% confidence intervals for the parameter β_2 in the predictive regression model (24). We consider rolling windows of 180 observations for the period 1950-2010. In the top line, we present the classic bootstrap (left panel) and the classic subsampling (right panel), while in the bottom line we consider our robust bootstrap (left panel) and our robust subsampling (right panel).




Figure 12: **Huber Weights under the Predictive Regression Model (21).** We plot the Huber weights for the predictive regression model (21) in the period 1980-2010.




Figure 13: **Huber Weights under the Predictive Regression Model (23).** We plot the Huber weights for the predictive regression model (23) in the period 1990-2010.




Figure 14: **Huber Weights under the Predictive Regression Model (24).** We plot the Huber weights for the predictive regression model (24) in the period 1950-2010.

$n = 120, b = 0.5$	0.9	0.95
Subsampling ($m = 10$)	[0.0417; 0.0833]	[0.0417; 0.0417]
Subsampling ($m = 20$)	[0.0833; 0.0833]	[0.0833; 0.0833]
Subsampling ($m = 30$)	[0.1250; 0.1250]	[0.1250; 0.1250]
Bootstrap ($m = 10$)	[0.0417; 0.3750]	[0.0417; 0.3333]
Bootstrap ($m = 20$)	[0.0833; 0.3333]	[0.0833; 0.3333]
Bootstrap ($m = 30$)	[0.1250; 0.3333]	[0.1250; 0.2500]

Table 1: **Subsampling and Block Bootstrap Lower and Upper Bounds for the Quantile Breakdown Point.** Breakdown point of the subsampling and the block bootstrap quantiles. The sample size is $n = 120$, and the block size is $m = 10, 20, 30$. We assume a statistic with breakdown point $b = 0.5$ and confidence levels $t = 0.9, 0.95$. Lower and upper bounds for quantile breakdown points are computed using Theorem 1.

$n = 120$	0.9	0.95
Subsampling ($m = 10$)	0.1750	0.1250
Subsampling ($m = 20$)	0.2500	0.2083
Subsampling ($m = 30$)	0.3250	0.2833
Bootstrap ($m = 10$)	0.5000	0.5000
Bootstrap ($m = 20$)	0.5000	0.5000
Bootstrap ($m = 30$)	0.4250	0.3583

Table 2: **Robust Subsampling and Robust Block Bootstrap for the studentized Statistic T_n .** Breakdown point of the robust subsampling and the robust block bootstrap quantiles for the studentized statistic T_n , in the predictive regression model (1)-(2). The sample size is $n = 120$, and the block size is $m = 10, 20, 30$. The quantile breakdown points are computed using Theorem 5.

	1980 – 1995	1985 – 2000	1990 – 2005	(1995 – 2010)
Bias-Corrected	0.0292 ^(**)	0.0167 ^(**)	0.0191 ^(**)	0.0156
Bonferroni	0.0236 ^(**)	0.0134 ^(**)	0.0117 ^(*)	0.0112
Bootstrap	0.0430 ^(**)	0.0175 ^(*)	0.0306 ^(**)	0.0355 ^(**)
Subsampling	0.0430 ^(**)	0.0175	0.0306 ^(**)	0.0355 ^(**)
R.Bootstrap	0.0405 ^(**)	0.0174 ^(**)	0.0245 ^(**)	0.0378 ^(**)
R.Subsampling	0.0405 ^(**)	0.0174 ^(**)	0.0245 ^(**)	0.0378 ^(**)

Table 3: **Point Estimates of Parameter β** . We report the point estimates of the parameter β in the predictive regression model (21) for the subperiods 1994-2009, 1980-1995, 1985-2000, 1990-2005 and 1995-2010, all consisting of 180 observations. In the second and third line we consider the bias-corrected method and the Bonferroni approach, respectively. In the fourth and fifth line we consider the standard bootstrap and standard subsampling, respectively. Finally, in the sixth and seventh line we consider our robust bootstrap and robust subsampling, respectively. (*) and (**) mean rejection at 10% and 5% significance level, respectively.

	1991 – 2006	1992 – 2007	1993 – 2008	1994 – 2009	1995 – 2010
Bootstrap	0.0357 ^(**)	0.0388 ^(**)	0.0435 ^(**)	0.0351 ^(*)	0.0406 ^(**)
Subsampling	0.0357 ^(**)	0.0388 ^(**)	0.0435 ^(**)	0.0351 ^(*)	0.0406 ^(**)
R.Bootstrap	0.0356 ^(**)	0.0387 ^(**)	0.0434 ^(**)	0.0348 ^(**)	0.0403 ^(**)
R.Subsampling	0.0356 ^(**)	0.0387 ^(**)	0.0434 ^(**)	0.0348 ^(**)	0.0403 ^(**)

	1991 – 2006	1992 – 2007	1993 – 2008	1994 – 2009	1995 – 2010
Bootstrap	0.3287 ^(**)	0.3203 ^(**)	0.3298 ^(**)	0.3011 ^(**)	0.3145 ^(**)
Subsampling	0.3287 ^(**)	0.3203 ^(**)	0.3298 ^(**)	0.3011 ^(**)	0.3145 ^(**)
R.Bootstrap	0.3368 ^(**)	0.3280 ^(**)	0.3381 ^(**)	0.3073 ^(**)	0.3214 ^(**)
R.Subsampling	0.3368 ^(**)	0.3280 ^(**)	0.3381 ^(**)	0.3073 ^(**)	0.3214 ^(**)

Table 4: **Point Estimates of Parameters β_1 and β_2** . We report the point estimates of parameters β_1 (first table) and β_2 (second table) in the predictive regression model (23) for the subperiods 1991-2006, 1992-2007, 1993-2008, 1994-2009 and 1995-2010, all consisting of 180 observations. In the second and third line we consider the standard bootstrap and standard subsampling, respectively. In the fourth and fifth line we consider our robust bootstrap and robust subsampling, respectively. (*) and (**) mean rejection at 10% and 5% significance level, respectively.

	1950 – 1995	1955 – 2000	1960 – 2005	1965 – 2010
Bootstrap	0.0724(**)	0.0301	0.0467(**)	0.0480(**)
Subsampling	0.0724(**)	0.0301	0.0467(**)	0.0480(*)
R.Bootstrap	0.0721(**)	0.0305(**)	0.0474(**)	0.0488(**)
R.Subsampling	0.0721(**)	0.0305(**)	0.0474(**)	0.0488(**)

	1950 – 1995	1955 – 2000	1960 – 2005	1965 – 2010
Bootstrap	−0.1509	−0.2926(**)	−0.2718(**)	−0.2187(*)
Subsampling	−0.1509	−0.2926(**)	−0.2718(**)	−0.2187
R.Bootstrap	−0.1532(**)	−0.2920(**)	−0.2701(**)	−0.2173(**)
R.Subsampling	−0.1532(**)	−0.2920(**)	−0.2701(**)	−0.2173(**)

Table 5: **Point Estimates of Parameters β_1 and β_2 .** We report the point estimates of parameters β_1 (rst table) and β_2 (second table) in the predictive regression model (24) for the subperiods 1950-1995, 1955-2000, 1960-2005 and 1965-2010, all consisting of 180 observations. In the second and third line we consider the standard bootstrap and standard subsampling, respectively. In the fourth and fifth line we consider our robust bootstrap and robust subsampling, respectively. (*) and (**) mean rejection at 10% and 5% significance level, respectively.