

Ordinal Efficiency, Fairness, and Incentives in Large Markets*

Qingmin Liu[†] and Marek Pycia[‡]

June 18, 2012

Abstract

Efficiency and symmetric treatment of agents are the primary goals of resource allocation in environments without transfers. Focusing on ordinal mechanisms in which no small group of agents can substantially change the allocation of others, we first show that uniform randomizations over deterministic efficient mechanisms are asymptotically ordinally efficient, that is, efficient *ex ante*. This implies that ordinal efficiency and *ex-post* Pareto efficiency become equivalent in large markets, and that many standard mechanisms are asymptotically ordinally efficient. Second, we show that all asymptotically ordinally efficient, symmetric, and asymptotically strategy-proof mechanisms lead to the same allocations in large markets.

*First draft: February 2011. The paper subsumes Pycia (2011b; 2011c). We thank Andrew Atkeson, Simon Board, Yeon-Koo Che, Federico Echenique, Hugo Hopenhayn, SangMok Lee, Moritz Meyer-ter-Vehn, George Mailath, Ichiro Obara, Joseph Ostroy, Mallesh Pai, Andy Postlewaite, Utku Ünver, Kyle Woodward, Leeat Yariv, William Zame, and seminar audiences at Northwestern Matching Workshop, UCLA, UPenn, SWET, and Caltech-UCLA Theory Conference for valuable comments. Keywords: ordinal efficiency, asymptotic ordinal efficiency, asymptotic strategy-proofness, symmetry, envy-freeness, *ex-post* Pareto efficiency, Random Priority, Probabilistic Serial, Randomized Hierarchical Exchange, Randomized Trading Cycles.

[†]Department of Economics, Columbia University, 1022 International Affairs Building, 420 West 118th Street New York, NY 10027; ql2177 (at) columbia.edu

[‡]Department of Economics, UCLA, 8283 Bunche, Los Angeles, CA 90095; pycia (at) ucla.edu; <http://pycia.bol.ucla.edu>

1 Introduction

Efficiency and symmetric treatment of participants (fairness) are the primary goals of allocation of resources in environments without monetary transfers such as assigning school seats to students and allocating university and public housing. In these examples – and in our model – there are many agents relative to the number of object types (also referred to as objects), each object type is represented by one or more indivisible copies, and each agent consumes at most one object copy.¹ Agents are indifferent among copies of the same object and have strict preferences among objects. Because object copies are indivisible, fair allocation mechanisms allocate objects randomly. Following the standard practice, we focus on mechanisms that are: (i) ordinal, that is, the random allocation depends only on agents' reports of their ordinal preferences over objects, and (ii) regular, that is, no small group of agents can substantially impact allocations of other agents.²

We know little about efficient and fair allocations in large finite markets without transfers. Prior literature established asymptotic efficiency of two mechanisms: Bogomolnaia and Moulin (2001) showed that an eating procedure they introduced and termed Probabilistic Serial is efficient, and Che and Kojima (2010) showed that a mechanism that randomly orders agents and then lets them pick objects in turn (Random Priority) is asymptotically equivalent to Probabilistic Serial, and thus also asymptotically efficient.

The present paper provides a criterion to establish asymptotic efficiency of mech-

¹See for instance Abdulkadiro Iu and Sönmez (2003) for school seat assignment and Chen and Sönmez (2002) for housing allocation. In school seat assignment, the number of students is large relative to the number of schools; in allocation of university housing, e.g. at Harvard, MIT, or UCLA, the set of rooms is partitioned into a small number of categories, and rooms in the same category are treated as identical.

²The literature focused on ordinal mechanisms because (a) real-life no-transfer mechanisms are typically ordinal, (b) learning and reporting one's preferences over sure outcomes is simpler than learning and reporting one's cardinal utilities, and (c) ordinal preferences over sure outcomes do not rely on agents' attitudes towards risk. For analysis that incorporates cardinal utilities see Hylland and Zeckhauser (1979) and Makowski, Ostroy, and Segal (1999). Regularity is a standard requirement in the study of large markets; see e.g. Champsaur and Laroque (1982).

anisms (Theorem 1), and shows that there is a unique way to achieve symmetric and asymptotically efficient allocations in an asymptotically strategy-proof way (Theorem 2). The first of these results implies that many known mechanisms are asymptotically efficient; in large markets there are no efficiency reasons to favor some of them over others. The second result establishes outcome equivalence of a large class of mechanisms, both known and unknown. The ongoing efforts to construct new mechanisms for large markets cannot succeed without relaxing some of the assumptions of Theorem 2.

Before discussing the results in more detail, let us review the standard concepts they rely on. The natural efficiency criterion in ordinal settings postulates that an allocation is efficient if no group of agents can advantageously swap equal-size probability shares in objects, and no agent can advantageously swap a probability share in an object for an equal-size share in an object that is unallocated with positive probability.³ We primarily study a natural asymptotic counterpart of this criterion: a mechanism is asymptotically efficient if the maximum size of advantageous swaps vanishes as the market becomes large. The baseline fairness criterion is symmetric, or equal, treatment of equal agents: an allocation is symmetric if any two agents who reported the same preference ranking are allocated objects according to the same distribution. A mechanism is strategy-proof if reporting preferences truthfully first-order stochastically dominates any other strategy. A mechanism is asymptotically strategy-proof if it is approximately strategy-proof, and the approximation error vanishes as the market becomes large.⁴

³Equivalently, an allocation is efficient if no other allocation first-order stochastically dominates it. This efficiency concept – known as ordinal efficiency or first-order-stochastic-dominance efficiency – was introduced by Bogomolnaia and Moulin (2001).

⁴ Strategy-proofness in ordinal settings has been studied by Gibbard (1978), Roth and Rothblum (1999), and Bogomolnaia and Moulin (2001). Asymptotic strategy-proofness is a weak incentive compatibility condition that has been intensively studied since – building on a seminal analysis of asymptotic incentives by Roberts and Postlewaite (1976) – Hammond (1979), Champsaur and Laroque (1982), and Jackson (1992) proved it for the Walrasian mechanism in large exchange economies. In our context, the case for restricting attention to strategy-proof mechanisms was made by Abdulkadiroğlu and Sönmez (2003) and Ergin and Sönmez (2006), while Azevedo and Budish (2011) review the literature on asymptotic strategy-proofness and make a general case in favor of

Our first main result, Theorem 1, establishes that uniform randomizations over Pareto-efficient deterministic mechanisms are asymptotically efficient. Thus, for instance, uniform randomizations over any Trading Cycle mechanisms of Pycia and Ünver (2009) are asymptotically efficient.⁵ As mentioned above, prior literature established asymptotic efficiency of Probabilistic Serial (Bogomolnaia and Moulin, 2001) and Random Priority (Che and Kojima, 2010).

One consequence of Theorem 1 is an equivalence of efficiency and ex-post Pareto efficiency in large markets. A random allocation is called ex-post Pareto efficient if it can be represented as a randomization over efficient deterministic allocations. Bogomolnaia and Moulin (2001) pointed out that ex post Pareto efficient mechanisms do not need to be efficient. It turns out that in large markets the two concepts become equivalent when we restrict attention to uniform randomizations.⁶

Our second main result, Theorem 2, establishes that all asymptotically efficient, symmetric, and asymptotically strategy-proof mechanisms are asymptotically equivalent that is the maximum difference in probability shares in an object an agent obtains under any two mechanisms vanishes as the market becomes large. Since it is straightforward to see that any uniform randomization over Trading Cycle mechanisms is regular, symmetric, and strategy-proof, Theorems 1 and 2 taken together imply that

nisms: to break away from the large market equivalence one needs to relax one of its assumptions. One important message from this theorem is that – in large markets – we cannot substantially improve upon the mechanisms we already know and use.⁷

While Theorem 2 likely is the first general result showing that all natural large market mechanisms coincide, the equivalence of some important special mechanisms has been proved before. Abdulkadiroğlu and Sönmez (1998) proved that Random Priority and the Core from Random Endowments (a uniform randomization over Gale’s Top Trading Cycles) are equivalent in environments in which each object has a single copy.⁸ In the large market setting of the present paper, Che and Kojima (2010) showed asymptotic equivalence of two mechanisms: Random Priority and Probabilistic Serial. Theorem 2 shows that the equivalence of these two mechanisms is not a coincidence but rather a fundamental property of allocation in large markets.

2 Model

A finite economy consists of a finite set of agents N , a finite set of objects Θ (also referred to as object types), and a finite set of object copies O . By $|a|$ we denote the number of copies of object $a \in \Theta$. To avoid trivialities, we assume that each object is represented by at least one copy.

Agents have unit demands and strict preferences over objects.⁹ We sometimes refer to an agent’s preference ranking as the type of the agent. \mathcal{P} denotes the set of preference rankings, and \mathcal{P}^N is the set of preference profiles. We assume that Θ contains the null object \emptyset (“outside option”), and we assume that it is not scarce, $|\emptyset| \geq |N|$. An object is called acceptable if it is preferred to \emptyset .

We can interchangeably talk about allocating objects and allocating copies; one natural interpretation is that object types represent schools, and object copies rep-

⁷For ongoing efforts to construct new mechanisms, see for instance Featherstone (2011).

⁸Carroll (2010) and Pathak and Sethuraman (2010) proved analogous results in environments with multiple copies.

⁹By \succsim we denote the weak-preference counterpart of a strict preference ranking \succ .

resent seats in these schools. We study random allocations. An *allocation* μ is given by probabilities $\mu(i, a) \in [0, 1]$ that agent i is assigned object a . An allocation is *deterministic* if $\mu(i, a) \in \{0, 1\}$ for all $i \in N$, $a \in \Theta$.¹⁰ All allocations studied in this paper are assumed to be feasible in the following sense

$$\begin{aligned} \sum_{i \in N} \mu(i, a) &\leq |a| \quad \text{for every } a \in \Theta, \\ \sum_{a \in \Theta} \mu(i, a) &= 1 \quad \text{for every } i \in N. \end{aligned}$$

The set of these random allocations is denoted by \mathcal{M} . A *mechanism* $\phi : \mathcal{P}^N \rightarrow \mathcal{M}$ is a mapping from the set of profiles of preferences over objects to the set of allocations.

We are primarily concerned with large but finite markets. To study them let us fix a sequence of finite economies $\langle N_q, \Theta, O_q \rangle_{q=1,2,\dots}$ in which the set of object types, Θ , is fixed while the set of agents N_q grows in q ; we assume throughout that $|N_q| \rightarrow \infty$ as $q \rightarrow \infty$. As discussed in the introduction, similar or more restrictive assumptions are standard in the study of large markets. To avoid repetition, in the sequel we refer to $\langle N_q, \Theta, O_q \rangle$ as the q -economy, and we assume that allocations μ_q and mechanisms ϕ_q are defined on q -economies. The number of copies of object a in the q -economy is denoted by $|a|_q$, and the set of random allocations in the q -economy is denoted by \mathcal{M}_q . Notice that we do not impose any assumptions on the sequence of sets of object copies, O_q ; in particular, we do not need to restrict attention to replica economies.¹¹

We study mechanisms ϕ_q in which the effect of reports of any small groups of agents other than j on the allocation of an agent j vanishes as $q \rightarrow \infty$. Formally, a sequence of mechanisms ϕ_q is *regular* if for every $\epsilon > 0$ there is $\delta > 0$ such that for

¹⁰A random allocation needs to be implemented as a lottery over deterministic allocations; Hylland and Zeckhauser (1979) and Bogomolnaia and Moulin (2001) showed how to implement random allocations. The implementation relies on Birkhoff and von Neumann's theorem.

¹¹Some of our results rely on additional assumptions on the ratio $|a|_q / |N_q|$, and we explicitly impose these assumptions when needed.

any preference profiles $\succ_{N_q}, \succ'_{N_q}$ and any agent $j \in N_q$ such that $\succ'_j = \succ_j$, if

$$\frac{|\{i \in N_q \mid \succ'_i \neq \succ_i\}|}{|N_q|} < \delta, \quad (1)$$

then

$$\max_{a \in \Theta} \left| \phi_q(\succ_{N_q})(j, a) - \phi_q(\succ'_{N_q})(j, a) \right| < \epsilon. \quad (2)$$

Regularity is a natural assumption; Champsaur and Laroque (1982) directly addresses the need for such an assumption. As the following remark illustrates, known mechanisms are regular.

Remark 1. Regularity of known mechanisms is straightforward to demonstrate. Take, for instance, Random Priority (Abdulkadiroğlu and Sönmez, 1998). To allocate objects, Random Priority first draws an ordering of agents from a uniform distribution over orderings, and then in turn allocates the first agent a copy of her most preferred object, allocates the second agent a copy of his most preferred object that still has unallocated copies, etc. This mechanism is regular provided for each object $a \in \Theta$ there is $\delta > 0$ such that

$$\liminf_{q \rightarrow \infty} |a|_q / N_q > \eta. \quad (3)$$

The proof fixes $\epsilon < \eta$ and has three steps.¹²

Step 1. Fix a priority ordering of agents. Conditional on this ordering, all allocations are deterministic. A change of preferences by ϵN_q agents can change the allocation of another agent j only if j takes one of the last ϵN_q copies of an object under at least one of the two preferences rankings submitted by the agents changing their preferences.

Step 2. The probability an agent takes one of the last ϵ copies of an object a under a preference profile vanishes as $q \rightarrow \infty$. Indeed, fix q and an ordering of agents other

¹²To see why assumption (3) is needed for regularity, consider an object $a \in \Theta$ and a sequence of economies such that $|a|_q = 1$. If the preference profile \succ_{N_q} is such that only two agents are interested in object a then a change of preferences by one of them has a large impact on the allocation of the other, violating regularity.

than $j \in N_q$, and consider probabilities conditional on such an ordering. If a is the favorite object for j then j would take it as long as it is available, and the conditional probability j takes one of the last ϵN_q copies of a is bounded above by $\frac{\epsilon N_q}{|a|_q - \epsilon N_q}$ (and for large q is bounded above by $\frac{\epsilon}{\eta - \epsilon}$). If there are objects ("better objects") that agent j prefers over a then j can take one of the last ϵN_q copies of a only after these better objects are exhausted; the probability of this happening is bounded above by $\frac{\epsilon}{\eta}$.

Step 3. By Step 2, the probability agent j takes one of the last ϵN_q copies of an object under one of two profiles of Step 1 is bounded above by $2|\Theta| \frac{\epsilon}{\eta - \epsilon}$ uniformly over agents and preference profiles. The regularity claim is thus true.

A similar argument shows that uniform randomizations over Hierarchical Exchange mechanisms of Pápai (2000) or Trading Cycles of Pycia and Ünver (2009) (extended to the setting with object copies by Pycia and Ünver (2011)) are regular provided the number of object copies satisfies condition (3).

3 Efficiency

The natural efficiency concept in our ordinal setting is the ordinal efficiency of Bogomolnaia and Moulin (2001): an allocation is efficient if agents cannot trade probability shares, and if no object is wasted.¹³ Formally, an allocation μ is *efficient* with respect to preference profile \succ iff (i) there is no cycle of agents i_0, i_1, \dots, i_n and objects a_0, a_1, \dots, a_n such that $\mu(i_k, a_k) > 0$ and $a_{k+1} \succ_{i_k} a_k$ (that is agents i_0, \dots, i_n cannot trade probability shares), and (ii) if $a \succ_i b$ and $\mu(i, b) > 0$, then all copies of a are allocated with probability 1 (that is no copy of a is wasted). This concept of efficiency naturally extends the concept of efficiency from deterministic allocations to random allocations. A mechanism ϕ is efficient if $\phi(\succ_N)$ is efficient for all preference profiles \succ_N .

¹³This concept is also known as first-order-stochastic-dominance efficiency because – in finite settings – efficiency of an allocation μ is equivalent to the lack of another allocation that first order stochastically dominates μ .

To define the asymptotic counterpart of this efficiency concept let us say that, for any $\epsilon > 0$, a random allocation μ is ϵ -efficient with respect to a preference profile \succ iff (i) there is no cycle of agents i_0, i_1, \dots, i_n and objects a_0, a_1, \dots, a_n such that $\mu(i_k, a_k) > \epsilon$ and $a_{k+1} \succ_{i_k} a_k$ (that is agents i_0, \dots, i_n cannot trade probability shares larger than ϵ), and (ii) if $a \succ_i b$ and $\mu(i, b) > \epsilon$, then all copies of a are allocated with probability at least $1 - \epsilon$ (that is no more than ϵ of a is wasted). Given a sequence of preference profiles \succ_{N_q} , a sequence of allocations μ_q is *asymptotically efficient* if for each $q = 1, 2, \dots$ there are positive $\epsilon(q) \xrightarrow{q \rightarrow \infty} 0$ such that μ_q is $\epsilon(q)$ -efficient with respect to \succ_{N_q} .

Which mechanisms are asymptotically efficient in large economies? We know some asymptotically efficient mechanisms: Bogomolnaia and Moulin (2001) constructed one such mechanism they called Probabilistic Serial. Recently, Che and Kojima (2010) showed that Random Priority is asymptotically equivalent to Probabilistic Serial; it is easy to see that their result implies that Random Priority is asymptotically efficient in the above sense.

The question remained open for other mechanisms. Many of the random mechanisms used in practice are obtained by uniformly randomizing over deterministic mechanisms. The main result of this section addresses this question for this large class of mechanisms. A mechanism $\phi : \mathcal{P}^N \rightarrow \mathcal{M}$ is a *uniform randomization* if there exists a mechanism $\psi : \mathcal{P}^N \rightarrow \mathcal{M}$ such that

$$\phi(\succ_{(1, \dots, |N|)})(i, a) = \sum_{\sigma: N \xrightarrow{1-1} N} \frac{1}{|N|!} \psi(\succ_{(\sigma(1), \dots, \sigma(|N|))})(\sigma(i), a)$$

In general, the component mechanism ψ can be random. We say that ψ is *ex-post Pareto efficient* if, for each preference profile \succ_N , the allocation $\psi(\succ_N)$ is equal to a mixture over efficient deterministic allocations. When ψ is ex-post Pareto efficient we say that ϕ is a *uniform randomization over efficient deterministic mechanisms*. A special case of interest is when ψ is an efficient deterministic mechanism. For

instance, if ψ is a serial dictatorship then ϕ is Random Priority. Other examples include randomizations over subsets of Papai’s Hierarchical Exchange or Pycia and Unver’s Trading Cycles.

Theorem 1. Regular sequences of uniform randomizations over efficient deterministic mechanisms are asymptotically efficient.

This result shows that all the above-mentioned mechanisms are asymptotically efficient. This result also shows that in large markets efficiency and ex-post Pareto efficiency become equivalent provided the randomization is uniform. This is in contrast to finite markets in which — following the pioneering work of Bogomolnaia and Moulin 2001 — it was well understood that ex-post Pareto efficiency is a weaker concept than efficiency.¹⁴

We sketch the proof below omitting some calculations provided in Appendix A. The key step in the proof is to use regularity to conclude that it is enough to prove the theorem for a carefully chosen subset of sequences of preference profiles \succ_q . We construct this set as follows. A sequence of preference-profiles \succ_{N_q} has *full support* if there exists $\delta > 0$ and \bar{q} such that for any $q > \bar{q}$, and for any ranking of objects $\succ \in \mathcal{P}$, the proportion of agents whose \succ_{N_q} -ranking agrees with \succ is above δ . Full support holds true uniformly on a class of sequences if they have full support with the same δ and \bar{q} . Full support implies that, as q grows, any preference ranking is represented by a non-vanishing fraction of agents.¹⁵

Let ϕ be a uniform randomization over Pareto-efficient deterministic mechanisms. Let us thus fix a sequence of preference profiles \succ_{N_q} ; regularity allows us to assume that this sequence has full support (see the first lemma in Appendix A for a detailed argument; this is the only use of regularity in the proof). To prove asymptotic

¹⁴See also McLennan (2002) and Abdulkadiro lu and Sönmez (2003).

¹⁵In a continuum economy, the counterpart of full support says that every ordering is represented with positive probability; in other words the distribution of orderings has full support. Our observations on full-support profiles remain valid if the assumption of non-vanishing representation is imposed only for ranking of objects \succ in which all non-null objects are acceptable.

ordinal efficiency we need to show that the allocations $\mu_q = \phi(\succ_{N_q})$ are $\epsilon(q)$ -ordinally efficient for some $\epsilon(q) \rightarrow 0$ as $q \rightarrow \infty$. By way of contradiction, assume that there is $\epsilon > 0$ and a sequence of $q_n \rightarrow \infty$ such that the allocations in the q_n -economies are not ϵ -efficient. For any $a \in \Theta$, symmetry of the uniform randomization implies that the probability $\mu_{q_n}(i, a)$ is the same for all agents i of a particular type $\succ \in \mathcal{P}$. Because there is a finite number of agent and object types, the compactness of $[0, 1]$ allows us also to further subsample the sequence q_n and assume that $\mu_{q_n}(i, a)$ converges to a constant $\mu_\infty(\succ, a) \in [0, 1]$.

We may assume that either condition (i) or condition (ii) fails for all q_n . Let us consider the case condition (i) fails for all q_n ; the proof in the remaining case follows the same steps. For each q_n there is a cycle of agents i_0, \dots, i_m and objects a_0, \dots, a_m such that i_k gets a higher-than- ϵ probability of a_k , and $a_{k+1} \succ_{i_k} a_k$ (subscripts modulo $m + 1$). Note we can have $m \leq |\mathcal{P}|$ for each q_n . By subsampling q_n we can get m and preference rankings \succ_{i_k} , $k = 1, \dots, m$ to be independent of q_n , and then $\mu_\infty(\succ_{i_k}, a_k) \geq \epsilon$.

By assumption there exist weights $\lambda_{k,q}$, $k = 1, \dots, K_q$ and Pareto-efficient deterministic mechanisms $\psi_{k,q}$ such that ϕ_q puts probability $\frac{\lambda_{k,q}}{|N_q|!}$ on $\psi_{k,q} \circ \sigma$ for all $k = 1, \dots, K_q$ and bijections $\sigma : N_q \rightarrow N_q$. Denote by M_q the random matrix that puts probability $\frac{\lambda_{k,q}}{|N_q|!}$ on the 0 – 1 matrix $\psi_{k,q} \circ \sigma(\succ_{N_q})$ indexed by $i \in N_q$ and $a \in \Theta$. Notice that the mean of $M_q(i, a)$ equals $\mu_q(i, a)$; the mean converges to $\mu_\infty(\succ_i, a)$ as $q \rightarrow \infty$.

A simple computation provided in the second lemma of Appendix A shows that for any agents i and j of the same preference type, and for any object a , the covariance between random variables $M_q(i, a)$ and $M_q(j, a)$ converges to zero as $q \rightarrow \infty$. We can thus apply the weak law of large numbers to random variables $M_q(i, a)$ and conclude that for any $\tilde{\epsilon} > 0$ and q large enough, the proportion of agents of type \succ_k is within $\tilde{\epsilon}$ of $\mu_\infty(\succ_k, a_k)$ with probability at least $1 - \tilde{\epsilon}$. This implies that there are some agents i_1, \dots, i_m of types \succ_1, \dots, \succ_m (respectively) who are allocated objects a_1, \dots, a_m (respectively) by some mechanism $\psi_{k,q} \circ \sigma$. This, however, contradicts the Pareto

efficiency of $\psi_{k,q}$ and concludes the proof.

4 Equivalence

As argued in the introduction, the natural postulates for allocation are efficiency, symmetry, and strategy-proofness. Unfortunately, we know that in finite markets no mechanism satisfies these three properties, see Bogomolnaia and Moulin (2001). Requiring only that the properties are satisfied in an asymptotic sense, we address the question of what mechanisms satisfy the three properties in large markets.

Let us first review the definitions of the standard concepts of symmetry and strategy-proofness. Symmetry is a basic fairness property of an allocation, and is also known as equal treatment of equals. Given preference profile \succ_N , a random allocation μ is *symmetric* if any two agents i and j who submitted the same ranking of objects, $\succ_i = \succ_j$, are allocated the same distributions over objects, $\mu(i, \cdot) = \mu(j, \cdot)$. A random mechanism ϕ is *strategy-proof* if for any agent $i \in N$ and any profile of preferences \succ_N , the allocation agent i obtains by reporting the truth, $\phi(\succ_i, \succ_{N-\{i\}})(i, \cdot)$, first-order stochastically dominates the allocation the agent can get by reporting another preference ranking \succ'_i , that is

$$\sum_{b \succsim_i a} \phi(\succ_i, \succ_{N-\{i\}})(i, b) \geq \sum_{b \succsim_i a} \phi(\succ'_i, \succ_{N-\{i\}})(i, b), \quad \forall a \in \Theta.$$

This is the standard concept of strategy-proofness introduced by Gibbard (1977). We further say that a random mechanism ϕ is ϵ -*strategy-proof* if for any agent $i \in N$ and any profile of preferences \succ_N ,

$$\sum_{b \succsim_i a} \phi(\succ_i, \succ_{N-\{i\}})(i, b) \geq \sum_{b \succsim_i a} \phi(\succ'_i, \succ_{N-\{i\}})(i, b) - \epsilon, \quad \forall a \in \Theta, \succ'_i \in \mathcal{P}.$$

A sequence of random mechanisms ϕ_q is asymptotically strategy-proof if for each

$q = 1, 2, \dots$ there are positive $\epsilon(q) \xrightarrow{q \rightarrow \infty} 0$ such that μ_q is $\epsilon(q)$ -strategy-proof. Footnote 4 discusses the literature on asymptotic strategy-proofness.

Theorem 2. *If two regular sequences of random mechanisms ϕ_q and ϕ'_q are asymptotically efficient, symmetric, and asymptotically strategy-proof, then they are asymptotically equivalent, that is*

$$\max_{\succ_{N_q} \in \mathcal{P}_q, i \in N_q, a \in \Theta} |\phi_q(\succ_{N_q})(i, a) - \phi'_q(\succ_{N_q})(i, a)| \rightarrow 0 \quad \text{as} \quad q \rightarrow \infty.$$

Theorem 2 has important ramifications for the ongoing work to construct improved allocation mechanisms: we cannot improve upon the allocations of mechanisms we already know without relaxing the requirement of (approximate) strategy-proofness or of symmetry. Featherstone (2011) provides an important example of such an ongoing work.¹⁶

Theorem 2 also immediately implies the following.

Corollary 1. *All asymptotically strategy-proof sequences of uniform randomizations over Pareto-efficient deterministic mechanisms are asymptotically equivalent.*

As a heuristic for why Theorem 2 is true let us consider a setting with a continuum of agents and assume that the mass of agents of type \succ , denoted $|N(\succ)|$, is strictly positive for all $\succ \in \mathcal{P}$. Let $|a|$ be the total mass of object a . One way to understand the continuum setting is as a limit of finite economies. This limit is well-defined provided we assume that (i) the sequence of preference profiles \succ_{N_q} is such that for each preference type $\succ \in \mathcal{P}$, the ratio $\frac{|\{i \in N_q | \succ_i = \succ\}|}{|N_q|}$ converges to a constant $|N(\succ)|$,

¹⁶The above results allow us to also obtain new insights into the strategy-proofness properties of Probabilistic Serial. Kojima and Manea (2010) showed that agents have incentives to report preferences truthfully in Probabilistic Serial if the number of copies is large enough relative to a measure of variability of an agent's utility, and Che and Kojima (2010) showed asymptotic strategy-proofness of Probabilistic Serial provided the number of copies has asymptotically the same rate of growth as $|N_q|$. Like Che and Kojima (2010), our results do not rely on assumptions on an agent's utility. The results allow us to relax Che and Kojima's assumption on the number of object copies, as well as to show that no assumption on the number of copies is needed for asymptotic strategy-proofness at full-support preference profile sequences.

and that $\frac{|a|_q}{|N_q|}$ converges to a positive constant $|a|$. Because of symmetry of ϕ and ϕ' , knowing the total mass of object a allocated to agents of type \succ is equivalent to knowing the probability an agent of type \succ obtains object a . We will refer to this probability as $\mu(i, a)$ or $\mu(\succ, a)$. A key role in our argument is played by the following strong fairness requirement introduced by Foley (1967).¹⁷ An allocation μ is *envy-free* if any agent i first-order stochastically prefers his allocation over the allocation of any other agent j , that is

$$\sum_{b \succsim_i a} \mu(i, b) \geq \sum_{b \succsim_j a} \mu(j, b), \quad \forall a \in \Theta.$$

Since $m(\succ) > 0$ for all types \succ , regularity, symmetry and asymptotic strategy-proofness of the sequence of mechanisms ϕ_q allow us to conclude that the limit allocation $(\mu(\succ, a))_{\succ \in \mathcal{P} \times \Theta}$ is envy-free. Thus, proving that there is a unique envy-free and efficient allocation is a step towards proving a weak version of Theorem 2 (weakened by the imposition of the additional restrictive assumptions listed above).¹⁸

To prove that efficiency and envy-freeness fully determine an allocation when $|N(\succ)| > 0$ for all $\succ \in \mathcal{P}$, first note that there are constants $t_a > 0$, $a \in \Theta$, such that $\mu(i, a) > 0$ imply $t_a = \sum_{b \succsim_i a} \mu(i, b)$. Indeed, take i, j such that $\mu(i, a), \mu(j, a) > 0$. Consider agent i' who ranks a first and otherwise ranks objects as agent i does; no envy and efficiency imply $\mu(i', a) = \sum_{b \succsim_i a} \mu(i, b)$. Similarly, for agent j' who ranks

¹⁷In the proof of Theorem 2 in the appendix we use asymptotic envy-freeness, a concept defined in Jackson and Kremer (2007) who noted its relation to incentive compatibility.

¹⁸The argument presented here can be used to prove a weak form of Theorem 2 in which we add assumption (b), and we restrict attention to convergence for preference profile sequences such that (a) obtains and $|N(\succ)| > 0$ for all $\succ \in \mathcal{P}$. Indeed, symmetry implies that $\phi_q(\succ_{N_q})(i, a)$ is fully determined by \succ_{N_q} , object a , and the preference type \succ of agent i ; we can thus write $\phi_q(\succ_{N_q})(\succ_i, a) = \phi_q(\succ_{N_q})(i, a)$. Full-support of \succ_{N_q} ensures that $\phi_q(\succ, a)$ is well-defined for any \succ and large q . At each preference profile along the sequence of profiles \succ_{N_q} , the mechanism is given by $(\phi(\succ_{N_q})(\succ, a))_{\succ \in \mathcal{P}, a \in \Theta} \in [0, 1]^{\mathcal{P} \times \Theta}$. Given compactness of this latter set, to show that two mechanisms ϕ_q and ϕ'_q coincide asymptotically along \succ_{N_q} it is enough to show that every convergent subsequence of $(\phi(\succ_{N_q})(\succ, a))_{\succ \in \mathcal{P}, a \in \Theta}$ converges to the same limit. We can thus consider a subsequence $(\phi(\succ_{N_q})(\succ, a))_{\succ \in \mathcal{P}, a \in \Theta}$ that converges to a limit $(\mu(\succ, a))_{\succ \in \mathcal{P} \times \Theta}$. To prove the weak version of the theorem it is then enough to show that $\mu(\succ, a)$ are uniquely determined by efficiency and envy-freeness.

a first and otherwise ranks objects as agent j does, $\mu(j', a) = \sum_{b \succsim_j a} \mu(j, b)$. Finally, no envy further implies that $\mu(i', a) = \mu(j', a)$, proving the claim. We refer to indices t_a as time and say that object a is not exhausted at all times $t < t_a$ and exhausted at all times $t \geq t_a$.¹⁹

Our assumptions uniquely determine agent-object pairs (i, a) such that $\mu(i, a) > 0$. To see this first note that efficiency and envy-freeness imply that agents i ranking a first get it with positive probability, $\mu(i, a) = t_a > 0$. No envy then implies that $\mu(i, a) > 0$ for all agents who rank a above all objects b that are not exhausted at time t_a . Let us denote the set of such types by $N(a)$. The definition of times t_{a_k} then implies that for $i \in N(a)$ we have

$$\mu(i, a) = t_a - t'_a \quad (4)$$

where t'_a equals 0 if agent i ranks a first and otherwise equals t_b for the object b that is \succsim_i -worst among objects \succsim_i ranks above a . Furthermore, only agents from $N(a)$ get a with positive probability. Indeed, if agent i ranks a below some object b non-exhausted at t_a , and $\mu(i, a) > 0$ then: (i) agent i gets object $b \succ a$ with probability below 1; (ii) consider agent j that ranks a first and b second and observe that agent j gets a with positive probability because $t_b > t_a > 0$. Equation (4) implies that j gets b with positive probability. Now (i) and (ii) imply that agents i and j have a profitable trade, a contradiction showing that $\mu(i, a) > 0$ iff $i \in N(a)$.

To conclude the proof let us rename the objects so that

$$0 < t_{a_1} \leq t_{a_2} \leq \dots \leq t_{a_{|\Theta|}}.$$

¹⁹This choice of terminology is motivated by parallels to the “eating-in-time” terminology of the Probabilistic Serial mechanism of Bogomolnaia and Moulin (2001).

The efficiency then implies that

$$t_{a_1} = \min \left\{ \frac{|a_1|}{|N(a_1)|}, 1 \right\}, \quad t_{a_2} = \min \left\{ t_{a_1} + \frac{|a_2| - t_{a_1} |\{i \in N(a_2) \mid i \text{ ranks } a_2 \text{ first}\}|}{|N(a_2)|}, 1 \right\},$$

and, proceeding in this way by induction, we can pin down all values t_a , thus uniquely determining allocation μ .

5 Comments on assumptions

We study regular mechanisms; the case for regularity has been made in the large market literature by, for instance, Champsaur and Laroque (1982). However, the gist of our results remain valid for all mechanisms, not only regular ones. If we drop the regularity assumption, the analogues of our results remain true for full-support sequences of preference profiles. In fact, the proofs of our results show that the results are true for full-support sequences of profiles whether or not the mechanism is regular, and we then employ regularity to extend the results to any preference profiles. Regularity is thus a dispensable assumption because full-support sequences are typical. Indeed, full-support sequences have asymptotically full measure in the following sense: a set \mathcal{S} of sequences of preference profiles is *asymptotically full-measure* if for every $\epsilon > 0$ there exists a sequence of sets $\mathcal{S}_q \subset \mathcal{P}_q$ of preference profiles in q -economies such that for q large enough the ratio $\frac{|\mathcal{S}_q|}{|\mathcal{P}_q|} > 1 - \epsilon$, and all sequences of profiles from \mathcal{S}_q are in the set of sequences \mathcal{S} .

Proposition 1. *Asymptotically full-support profiles have asymptotically full-measure.*

Proof. Let $\mathcal{S}_q^\delta \subset \mathcal{P}_q$ be the set of preference profiles such that, for any ranking of objects \succ in the q -economy, the proportion of agents whose ranking agrees with \succ to $|N_q|$ is above δ . Take $\epsilon > 0$ and notice that for q large enough there exists $\delta(\epsilon) > 0$ such that $\frac{|\mathcal{S}_q^{\delta(\epsilon)}|}{|\mathcal{P}_q|} > 1 - \epsilon$. To complete the proof it is enough to set $\mathcal{S}_q = \mathcal{S}_q^{\delta(\epsilon)}$. \square

Another assumption we can relax is symmetry. While the heuristic argument for

Theorem 2 made substantial use of symmetry, the formal argument from Appendix B does not, and this argument remains valid when Theorem 2 is strengthened by relaxing the symmetry assumption to its asymptotic counterpart.²⁰

6 Conclusion

Theorem 1 shows that – in ordinal settings – many known mechanisms are asymptotically efficient, and Theorem 2 establishes that effectively there is only one way to allocate objects in an efficient, symmetric, and strategy-proof way. Thus, in large markets, the choice among ordinal mechanisms needs to be based on criteria other than efficiency or fairness. This has important implications for ongoing efforts to construct new ordinal mechanisms.

In the current paper, we study the canonical single-unit assignment model. In Liu and Pycia [2011], we extended the results to multi-unit environments.²¹

A Proof of Theorem 1

The only elements missing in the proof of Theorem 1 in the main text are the following two lemmas:

Lemma 1. *If ϕ is regular and $\phi(\succ_{N_q})$ is asymptotically efficient for full-support sequences $(\succ_{N_q})_{q=1,2,\dots}$, then ϕ is asymptotically efficient.*

Proof. Fix $\delta > 0$ and a large positive integer \bar{q} , and let $\mathcal{S}_{\delta,q}$ be the class of sequences of preference profiles \succ_{N_q} such that for all $q > \bar{q}$, each ranking $\succ \in \mathcal{P}$ is

²⁰Given a sequence of preference profiles \succ_{N_q} , a sequence of random allocations μ_q is asymptotically symmetric if

$$\max_{i,j \in N_q \text{ such that } \succ_i = \succ_j, a \in \Theta} |\mu_q(i, a) - \mu_q(j, a)| \rightarrow 0 \text{ as } q \rightarrow \infty.$$

A mechanism is *asymptotically symmetric* if the convergence is uniform across preference profiles.

²¹Cf Kojima (2009), Budish, Che, Kojima, and Milgrom (2011), and Pycia (2011a) for models of multiple unit assignment. In a follow up work, Heo (2011) extended Lemma 3 to the non-strict preference environment of Katta and Sethuraman (2006).

represented in \succ_{N_q} by at least a fraction δ of agents. The asymptotic efficiency of $\phi_q(\succ_{N_q})$ obtains uniformly on $\mathcal{S}_{\delta,q}$. Indeed, if not then there would exist $\epsilon > 0$ and a sequence of sequences of preference profiles $(\succ_{N_q}^k)_{q=1,2,\dots}$ indexed by $k = 1, 2, \dots$ such that no $\phi_q(\succ_{N_q}^q)$ is ϵ -ordinally efficient; a contradiction because the sequence of profiles $(\succ_{N_q}^q)_{q=1,2,\dots}$ belongs to $\mathcal{S}_{\delta,q}$ and in particular has full-support.

To finish the proof, take any sequence of profiles $\succ_{N_q} \in \mathcal{P}^{N_q}$. There is a sequence of profiles $(\succ'_{N_q})_{q=1,2,\dots} \in \mathcal{S}_{\delta,q}$ such that (1) is satisfied. The uniform asymptotic efficiency on $\mathcal{S}_{\delta,q}$ and inequality (2) imply that $\phi_q(\succ_{N_q})$ satisfies conditions (i)-(ii) of ϵ -efficiency uniformly on $\succ_{N_q} \in \mathcal{P}^{N_q}$, for large q (the ϵ in ϵ -efficiency needs to be slightly larger than ϵ in (2), e.g. twice the size). By taking $\epsilon \rightarrow 0$ we prove the claim. QED

Lemma 2. Let $\lambda_{k,q} > 0$, $k = 1, \dots, K_q$, add-up to 1, let $\psi_{k,q}$ be Pareto-efficient deterministic mechanisms, and let M_q put probability $\frac{\lambda_{k,q}}{|N_q|!}$ on $\psi_{k,q} \circ \sigma$ for all $k = 1, \dots, K_q$ and bijections $\sigma : N_q \rightarrow N_q$. Then, for any agents i and j of the same preference type, and for any object a , the correlation between random variables $M_q(i, a)$ and $M_q(j, a)$ converges to zero as $q \rightarrow \infty$. The convergence is uniform on any class of uniformly asymptotically full-support profiles.

Proof. Fix q , a preference profile in \mathcal{P}^{N_q} , and an ordering $\succ \in \mathcal{P}$. Denote by n_a the number of agents of type \succ getting a under the fixed preference profile, and let $n = \sum_{a \in \Theta} n_a$. The covariance between two agents i and j of type \succ getting a (that is, the covariance between $M_q(i, a)$ and $M_q(j, a)$) is the average of such covariances conditional on the profile of numbers n_a , $a \in \Theta$. The symmetry among agents of the same preference type implies that conditional on a profile of n_a , $a \in \Theta$, the covariance between $M_q(i, a)$ and $M_q(j, a)$ is

$$\begin{aligned} & \frac{n_a}{n} \frac{n_a - 1}{n - 1} \left(1 - \frac{n_a}{n}\right)^2 + 2 \frac{n_a}{n} \left(\frac{n - n_a}{n - 1}\right) \left(1 - \frac{n_a}{n}\right) \left(0 - \frac{n_a}{n}\right) + \\ & + \left(\frac{n - n_a}{n}\right) \left(\frac{n - 1 - n_a}{n - 1}\right) \left(0 - \frac{n_a}{n}\right)^2 = \frac{n_a(n_a - n)}{n^2(n - 1)} \in \left[-\frac{1}{4(n - 1)}, 0\right]. \end{aligned}$$

The covariance thus converges to 0 as $q \rightarrow \infty$ because n , the number of agents of type \succ , grows to infinity along asymptotically full-support profiles. The uniform convergence claim is straightforward. QED

B Proof of Theorem 2

In the proof we will use the ingenious construction of the Probabilistic Serial mechanism of Bogomolnaia and Moulin (2001); this mechanism is both efficient and envy-free. Probabilistic Serial treats copies of an object type as a pool of probability shares of the object type. Given preference profile \succ_N , the random allocation produced by Probabilistic Serial can be determined through an “eating” procedure in which each agent “eats” probability shares of the best acceptable and available object with speed 1 at every time $t \in [0, 1]$; an object a is available at time t if its initial endowment $\theta^{-1}(a)$ is larger than the sum of shares that have been eaten by time t .

Formally, at time $t = 0$, the total quantity of available shares of object type $a \in \Theta$ is $Q_a(0) = |\theta^{-1}(a)|$, and for times $t \in [0, 1)$ we define the set of available objects $A(t) \subseteq \Theta$ and the available quantity $Q_a(t)$ of probability shares of object $a \in \Theta$ through the following system of integral equations

$$\begin{aligned} A(t) &= \{a \in \Theta \mid Q_a(t) > 0\}, \\ Q_a(t) &= Q_a(0) - \int_0^t |\{i \in N \mid a \in A(\tau) \text{ and } \forall b \in A(\tau) \ a \succsim_i b\}| d\tau. \end{aligned}$$

We say that *agent i eats from object a at time t* iff $a \in A(t)$ and $\forall b \in A(t)$, $a \succsim_i b$. If stopped at time t , the eating procedure allocates object $a \in \Theta$ to agent $i \in N$ with probability

$$\psi^t(i, a) = \int_0^t \chi(i \text{ eats from } a \text{ at time } \tau) d\tau,$$

where the Boolean function $\chi(\text{statement})$ takes value 1 if the statement is true and 0 otherwise. The allocation $\psi(i, a)$ of *Probabilistic Serial* is given by the eating

procedure stopped at time 1; that is $\psi = \psi^1$.

The continuity of the functions Q_a implies that for any time $T \in [0, 1)$ and any $\eta > 0$ sufficiently small, any agent i eats the same object for all $t \in [T, T + \eta)$. In the eating procedure there are some critical times when one or more objects get exhausted. At this time some of the available quantity functions Q_a have kinks; at other times their slope is constant.²²

Before proving Theorem 2, let us also define asymptotic envy-freeness. First, given an $\epsilon > 0$ and preference profile \succ_{N_q} , let us say that an allocation μ is ϵ -envy free if

$$\sum_{b \succsim_i a} \mu(i, b) + \epsilon \geq \sum_{b \succsim_i a} \mu(j, b), \quad \forall a \in \Theta, \forall i, j \in N_q.$$

Given a sequence of preference profiles \succ_{N_q} , we say that a sequence of allocations μ_q is asymptotically envy free if there are positive $\epsilon_q \rightarrow 0$ such that μ_q is ϵ_q -envy free. We say that a sequence of mechanisms ϕ_q is asymptotically envy-free if there are positive $\epsilon(q) \xrightarrow{q \rightarrow \infty} 0$ such that allocations $\phi_q(\succ_{N_q})$ are ϵ_q -envy free for all \succ_{N_q} .

Lemma 3. Fix a full-support sequence of preference profiles. If a sequence of random allocations μ_q is asymptotically efficient and asymptotically envy-free, then it coincides asymptotically with the allocations of Probabilistic Serial. Moreover, if a sequence of mechanisms ϕ_q is asymptotically efficient, symmetric, and asymptotically envy-free, and a class of profile sequences \mathcal{S} has uniformly full support, then ϕ_q converges to Probabilistic Serial uniformly on this class, that is

$$\max_{(\succ_{N_q})_{q=1,2,\dots}} \max_{i \in \mathcal{S}, l \in N_q, a \in \Theta} |\phi_q(\succ_{N_q})(i, a) - \phi'_q(\succ_{N_q})(i, a)| \rightarrow 0 \quad \text{as } q \rightarrow \infty.$$

Proof. Fix any sequence of full-support preference profiles \succ_{N_q} and an asymptotically efficient and asymptotically envy-free sequence of allocations μ_q . Asymptotic

²²This structure of quantity functions Q_a implies that we can define the allocation of Probabilistic Serial through a system of difference equations; such definitions are given in Bogomolnaia and Moulin (2001), and, for the environment with copies, in Kojima and Manea (2010).

efficiency implies that for any small $\epsilon > 0$ and large $M > 0$ there is \bar{q} such that for $q \geq \bar{q}$, the allocations are $\frac{\epsilon}{M^2}$ -efficient, and, in particular, there are no two agents who could swap probability shares of size $\frac{\epsilon}{M^2}$ in some two objects. By asymptotic envy-freeness we can assume that for $q \geq \bar{q}$ each agent i 's allocation $\frac{\epsilon}{M^2}$ -first order stochastically dominates allocations of any other agent j in agent i 's preferences, that is

$$\sum_{b \succsim_i a} \mu_q(i, b) - \sum_{b \succsim_i a} \mu_q(j, b) \geq -\frac{\epsilon}{M^2} \quad \text{for all } a \in \Theta.$$

We fix $q \geq \bar{q}$ and, to economize on notation, we drop the q -subscript when referring to this fixed economy $N = N_q$ and its allocation $\mu = \mu_q$. We also drop the preference argument when referring to the allocation of Probabilistic Serial ψ .

To prove the lemma it is enough to show that

$$\sum_{a' \succsim_i a} \mu(i, a') \geq \sum_{a' \succsim_i a} \psi^t(i, a') - \frac{\epsilon}{M} \tag{5}$$

for all $t \in [0, 1]$, agents i , and objects $a \in \Theta$. Indeed, this set of inequalities for $t = 1$, together with efficiency of the Probabilistic Serial ψ^1 , imply that $|\mu(i, a) - \psi^1(i, a)| < \epsilon$ for all i and a , provided $M \geq |\Theta|$.

By way of contradiction, assume the above inequality fails for some time, agent, and object. Let T be the infimum of $t \in [0, 1]$ such that there exists $i \in N$ and $b \in \Theta$ such that $\sum_{a' \succsim_i b} \mu(i, a') < \sum_{a' \succsim_i b} \psi^t(i, a') - \frac{\epsilon}{M}$. Since there are a finite number of agents and objects, there is an agent and object for which the infimum is realized; let us fix such an agent and such an object and call them i and b , respectively. Let us assume that b is the highest ranked object in i 's preferences for which the infimum is realized.

Step 1. Inequalities (5) are satisfied at $t = T$ because the mapping $t \mapsto \psi^t(i, a)$ is continuous and the inequalities are satisfied for $t \in [0, T)$. In particular, the cutoff time T belongs to $[0, 1)$.

Step 2. At time T of the eating procedure, agent i must be eating from b . Indeed,

if i is eating from an object $a \succ_i b$ at T , then $\psi^T(a') = 0$ for all objects $a' \prec_i a$, and hence if (5) is violated for agent i and object b then it is violated for agent i and object a . This would contradict the assumption that i ranks b above all other objects for which the infimum T is realized. If i is eating from an object $a \prec_i b$ at time T then $\sum_{a' \succsim_i b} \mu(i, a') \geq \sum_{a' \succsim_i b} \psi^T(i, a') - \frac{\epsilon}{M} = \sum_{a' \succsim_i b} \psi^t(i, a') - \frac{\epsilon}{M}$ for t just above T , again contrary to T being the infimum of t at which (5) is violated for i and b .

Step 3. Agent i gets object b or better with probability $T - \frac{\epsilon}{M}$, that is $\sum_{a' \succsim_i b} \mu(i, a') = T - \frac{\epsilon}{M}$. Indeed, by Step 2, agent i is eating from b at time T in the eating procedure, and thus $\sum_{a' \succsim_i b} \psi^T(i, a') = T$. Because (5) is satisfied for $t = T$, we get $\sum_{a' \succsim_i b} \mu(i, a') \geq \sum_{a' \succsim_i b} \psi^T(i, a') - \frac{\epsilon}{M} = T - \frac{\epsilon}{M}$. The inequality is binding because functions $t \mapsto \psi^t(i, a')$ are continuous in t and T is the infimum of times at which (5) is violated.

Step 4. If b is the favorite object of agent $j \in N$, then $\mu(j, b) \in [T - \frac{\epsilon}{M}, T - \frac{\epsilon}{M} + \frac{\epsilon}{M^2}]$. Indeed, by Step 1, the top choice object b is still available at time t in the eating procedure, and thus $\psi^T(j, b) = T$. Because (5) is satisfied at time T we thus get $\mu(j, b) \geq T - \frac{\epsilon}{M}$. Furthermore, envy-freeness of μ implies that $\mu(j, b) \leq T - \frac{\epsilon}{M} + \frac{\epsilon}{M^2}$ as otherwise the outcome of agent i would not $\frac{\epsilon}{M^2}$ -first-order stochastically dominate for agent i the outcome of agent j .

Step 5. If b is the favorite object of agent $j \in N$, then $\psi^1(j, b) > T$. Indeed, if not, then in the eating procedure b would be exhausted at time T , contrary to i eating b at time T and thus at some times $t > T$.

Step 6. There is an agent $k \in N$ such that $\mu(k, b) > \psi^1(k, b) + \frac{3\epsilon}{M^2}$. Indeed, by the asymptotic full-support assumption at least a fraction δ of agents ranks b as their first choice. Steps 3 and 4 imply that under μ these agents get at least $\frac{(M-1)\epsilon}{M^2}$ less b than they get under ψ . Because $\delta > 0$ and is independent of M , for M large enough the $\frac{\epsilon}{M^2}$ -efficiency of μ implies that there must be another agent k who gets $\frac{3\epsilon}{M^2}$ more b under μ than under ψ^1 .

Step 7. There is an object $c \neq b$ that agent k from Step 6 ranks just above b .

Indeed, the claim follows from Steps 4, 5, and 6.

Let us fix agent k and object c satisfying Steps 6 and 7.

Step 8. Under μ , agent k gets object b or better with probability strictly higher than $T - \frac{\epsilon}{M} + \frac{3\epsilon}{M^2}$. Indeed, Step 1 and the availability of object b at time T in the eating procedure imply that $\sum_{a \succsim_k c} \mu(k, a) \geq \sum_{a \succsim_k c} \psi^T(k, a) - \frac{\epsilon}{M} = T - \psi^T(k, b) - \frac{\epsilon}{M} \geq T - \psi^1(k, b) - \frac{\epsilon}{M}$. The claim then follows from Step 6.

To conclude the proof, notice that by the asymptotic full support assumption, there exists an agent j who ranks objects in the same way as agent k except that j puts b first. By Step 6, $\mu(k, b) > \frac{\epsilon}{M^2}$, and thus the lack of swaps of size $\frac{\epsilon}{M^2}$ (the consequence of $\frac{\epsilon}{M^2}$ -efficiency of μ) implies that $\sum_{a \succ_k b} \mu(j, a) < \frac{\epsilon}{M^2}$. Step 4 thus implies that under μ the probability j gets object c or better is between $T - \frac{\epsilon}{M}$ and $T - \frac{\epsilon}{M} + \frac{2\epsilon}{M^2}$, and, by Step 8, it is smaller than the probability k gets these objects. This contradicts envy-freeness of μ . The contradiction proves (5), and the first part of the lemma.

An examination of the above argument shows that the choice of ϵ , M , and \bar{q} can be made uniformly for $\mu_q = \phi_q(\succ_{N_q})$ on a class of preference profile sequences with uniformly full-support, proving the second part of the lemma.

Proof of Theorem 2. First note that it is enough to prove the result assuming that ϕ'_q are Probabilistic Serial. Second, notice that regularity, symmetry, and asymptotic strategy-proofness imply that $\phi_q(\succ_{N_q})$ is asymptotically envy free for all \succ_{N_q} . The above lemma shows a uniform convergence of $\phi(\succ_{N_q})$ to Probabilistic Serial for all sequences of preference profiles \succ_{N_q} such that for some $\delta > 0$ and positive integer \bar{q} , for all $q > \bar{q}$ each ranking $\succ \in \mathcal{P}$ is represented in \succ_{N_q} by at least fraction δ of agents. Let us then fix any sequence of profiles \succ_{N_q} , and derive the convergence for \succ_{N_q} from the regularity of ϕ_q and the convergence for asymptotically full-support sequences of profiles \succ'_{N_q} such that (1) is satisfied. The convergence is uniform across preference profiles, proving Theorem 2. QED

References

- ABDULKADIROĞLU, A. AND T. SÖNMEZ (1998): “Random Serial Dictatorship and the Core from Random Endowments in House Allocation Problems,” *Econometrica*, 66, 689–701.
- ABDULKADIROĞLU, A. AND T. SÖNMEZ (2003): “Ordinal Efficiency and Dominated Sets of Assignments,” *Journal of Economic Theory*, 112, 157–172.
- ABDULKADIROĞLU, A. AND T. SÖNMEZ (2003): “School Choice: A Mechanism Design Approach,” *American Economic Review*, 93, 729–747.
- AZEVEDO, E. AND E. BUDISH (2011): “Strategyproofness for “Price-Takers” as a Desideratum for Market Design,” Unpublished manuscript.
- BOGOMOLNAIA, A. AND H. MOULIN (2001): “A New Solution to the Random Assignment Problem,” *Journal of Economic Theory*, 100, 295–328.
- BUDISH, E., Y.-K. CHE, F. KOJIMA, AND P. MILGROM (2011): “Implementing Random Assignments: A Generalization of the Birkhoff - von Neumann Theorem,” University of Chicago, Columbia University, Stanford University, unpublished mimeo.
- CARROLL, G. (2010): “A General Equivalence Theorem for Allocation of Indivisible Objects,” MIT, Working paper.
- CHAMPSAUR, P. AND G. LAROQUE (1982): “A Note on Incentives in Large Economies,” *The Review of Economic Studies*, 49, 627–635.
- CHE, Y.-K. AND F. KOJIMA (2010): “Asymptotic Equivalence of Random Priority and Probabilistic Serial Mechanisms,” *Econometrica*, 78, 1625–1672.
- CHEN, Y. AND T. SÖNMEZ (2002): “Improving Efficiency of On-campus Housing: An Experimental Study,” *American Economic Review*, 92, 1669–1686.
- ERGIN, H. AND T. SÖNMEZ (2006): “Games of School Choice under the Boston Mechanism,” *Journal of Public Economics*, 90, 215–237.
- FEATHERSTONE, C. R. (2011): “Rank Efficiency: Investigating a Widespread Ordinal Welfare Criterion,” .
- FOLEY, D. (1967): “Resource Allocation and the Public Sector,” *Yale Economic Essays*, 7, 45–98.
- GIBBARD, A. (1977): “Manipulation of Schemes That Mix Voting with Chance,” *Econometrica*, 45, 665–681.
- (1978): “Straightforwardness of Game Forms with Lotteries as Outcomes,” *Econometrica*, 46, 595–614.
- HAMMOND, P. J. (1979): “Straightforward Individual Incentive Compatibility in Large Economies,” *The Review of Economic Studies*, 46, 263–282.

- HYLLAND, A. AND R. ZECKHAUSER (1979): “The Efficient Allocation of Individuals to Positions,” *Journal of Political Economy*, 87, 293–314.
- JACKSON, M. O. (1992): “Incentive Compatibility and Economic Allocation,” *Economic Letters*, 299–302.
- JACKSON, M. O. AND I. KREMER (2007): “Envy-freeness and Implementation in Large Economies,” *Review of Economic Design*, 11, 185–198.
- KATTA, A.-K. AND J. SETHURAMAN (2006): “A Solution to The Random Assignment Problem on The Full Preference Domain,” *Journal of Economic Theory*, 131, 231–250.
- KOJIMA, F. (2009): “Random Assignment of Multiple Indivisible Objects,” *Mathematical Social Sciences*, 57, 134–142.
- KOJIMA, F. AND M. MANEA (2010): “Incentives in the Probabilistic Serial Mechanism,” *Journal of Economic Theory*, 144, 106–123.
- MAKOWSKI, L., J. M. OSTROY, AND U. SEGAL (1999): “Efficient Incentive Compatible Economies Are Perfectly Competitive,” *Journal of Economic Theory*, 85, 169–225.
- MANEA, M. (2009): “Asymptotic Ordinal Inefficiency of Random Serial Dictatorship,” *Theoretical Economics*, 4, 165–197.
- MCLENNAN, A. (2002): “Ordinal Efficiency and The Polyhedral Separating Hyperplane Theorem,” *Journal of Economic Theory*, 105, 435–449.
- MIRALLES, A. (2008): “School Choice: The Case for the Boston Mechanism,” Boston University, unpublished mimeo.
- PÁPAI, S. (2000): “Strategyproof Assignment by Hierarchical Exchange,” *Econometrica*, 68, 1403–1433.
- PATHAK, P. A. AND J. SETHURAMAN (2010): “Lotteries in Student Assignment: An Equivalence Result,” *Theoretical Economics*, forthcoming.
- PYCIA, M. (2011a): “Assignment with Multiple-Unit Demand and Responsive Preferences,” .
- (2011b): “On Envy-Freeness,” .
- (2011c): “Ordinally Efficient Allocation,” .
- PYCIA, M. AND M. U. ÜNVER (2009): “Incentive Compatible Allocation and Exchange of Discrete Resources,” UCLA and Boston College, unpublished mimeo.
- PYCIA, M. AND U. ÜNVER (2011): “Trading Cycles for School Choice,” .
- ROBERTS, D. J. AND A. POSTLEWAITE (1976): “The Incentives for Price-Taking Behavior in Large Exchange Economies,” *Econometrica*, 44, 115–127.

ROTH, A. E. AND U. ROTHBLUM (1999): "Truncation Strategies in Matching Markets: In Search of Advice for Participants," *Econometrica*